

On centralizers of parabolic subgroups in Coxeter groups

Koji Nuida

September 2, 2009

Abstract

Let W be an arbitrary Coxeter group, possibly of infinite rank. We describe a decomposition of the centralizer $Z_W(W_I)$ of an arbitrary parabolic subgroup W_I into the center of W_I , a Coxeter group and a subgroup defined by a 2-cell complex. Only information about finite parabolic subgroups is required in an explicit computation. Moreover, by using our description of $Z_W(W_I)$, we reveal a further strong property of the action of the third factor on the second factor, in particular on the finite irreducible components of the second factor.

1 Introduction

1.1 Background and summary of this work

A group W is called a Coxeter group if it has a generating set S such that W admits a presentation of the following form:

$$W = \langle S \mid (st)^{m_{s,t}} = 1 \text{ for all } s, t \in S \text{ such that } m_{s,t} < \infty \rangle ,$$

where the data $m_{s,t} \in \{1, 2, \dots\} \cup \{\infty\}$ are symmetric in $s, t \in S$ and we have $m_{s,t} = 1$ if and only if $s = t$. The pair (W, S) of a group W and such a generating set S is called a Coxeter system. Coxeter groups and some associated objects, such as root systems, appear frequently in various topics of mathematics such as geometry, representation theory, combinatorics, etc. In connection with such related topics, algebraic or combinatorial properties of Coxeter systems and those associated objects have been well studied, forming a long history and establishing many beautiful theories (see e.g., [16] and references therein). However, group structures of Coxeter groups themselves, especially those of infinite Coxeter groups, have not been studied so well; for example, even a fundamental property that an infinite irreducible Coxeter group is always directly indecomposable as an abstract group had not been known until very recent works [21, 26].

The object of this paper is the centralizer $Z_W(W_I)$ of any parabolic subgroup, which is the subgroup $W_I = \langle I \rangle$ generated by a subset I of S , in an arbitrary Coxeter group W . We emphasize that we do not place any restriction on W or W_I such as finiteness of cardinalities or finiteness of ranks. Analogously to other classes of groups such as finite simple groups, it is natural to hope that centralizers of some typical subgroups of Coxeter groups are useful to investigate the structural properties of Coxeter groups as abstract groups. Indeed, the results of this paper are applied in an indispensable manner to the author's recent study on the isomorphism problem in Coxeter groups (see [18]). See Section 1.2 for other applications.

We give a brief summary of the results of this paper. We present a decomposition of the centralizer $Z_W(W_I)$ in the following manner:

$$Z_W(W_I) = Z(W_I) \times (W^{\perp I} \rtimes B_I) .$$

Here the first factor $Z(W_I)$ is the center of W_I whose structure is well known; the second factor $W^{\perp I}$ is the subgroup generated by the reflections along the roots orthogonal to all roots associated to the subsystem (W_I, I) of (W, S) ; and the third factor B_I is a certain subgroup. For the second factor $W^{\perp I}$, an existing general theorem implies that such a reflection subgroup $W^{\perp I}$ forms a Coxeter group with canonical generating set; in this paper we give an explicit description of the canonical generating set and the corresponding fundamental relations of $W^{\perp I}$ by using some graphs and some 2-cell complex. The conjugation action of the third factor B_I on $W^{\perp I}$ induces automorphisms on $W^{\perp I}$ as a Coxeter system, i.e., those leaving the canonical generating set invariant. We describe the subgroup B_I as an extension of the fundamental group Y_I of the above-mentioned 2-cell complex by certain symmetries, and determine the action of B_I in terms of this complex.

One may feel that the above-mentioned result is similar to the preceding result on the normalizer $N_W(W_I)$ of W_I given by Brigitte Brink and Robert B. Howlett [5] (or by Howlett [14] for the case of a finite W). In fact, our decomposition of the centralizer looks like the decomposition of the normalizer given by their work, and some properties of our decomposition are derived from the corresponding properties for normalizers. However, it is worthy to emphasize that some fundamental properties of our decomposition are essentially new and desirable, and cannot be derived from the result on normalizers. Moreover, those new properties enable us to prove (under a certain assumption on I) the following strong property, which is another main result of this paper, of the centralizer $Z_W(W_I)$ that has been known for neither the centralizers nor the normalizers: The above-mentioned subgroup Y_I of $Z_W(W_I)$ (which occupies a fairly large part of the third factor B_I) acts *trivially* on every finite irreducible component of the Coxeter group $W^{\perp I}$. (This novel property plays a central role in the author's recent study on the isomorphism problem in Coxeter groups; see [18].) Thus the result of this paper is not just an analogy of the preceding result, but gives further novel insights into the structural properties of Coxeter groups. See Section 1.2 for detailed explanations. Conversely, as a by-product of our result, we also give another decomposition of $N_W(W_I)$ which shares the above-mentioned desirable properties with the decomposition of $Z_W(W_I)$.

1.2 Related works

As mentioned in Section 1.1, a similar problem of describing the normalizer $N_W(W_I)$ of a parabolic subgroup W_I was studied for finite W by Howlett [14] and for general W by Brink and Howlett [5]. Some part of our result is closely related to their preceding result, however some part of our result is essentially new and not derived from their work. More precisely, in our description of the centralizer $Z_W(W_I)$ we use a groupoid C , which turns out to be a covering of the groupoid G developed in their work, therefore some (but not all) properties of C are inherited from those of G . Our groupoid C has $W^{\perp I} \rtimes Y_I$ as a vertex group, and by analyzing the groupoid we give a presentation of $W^{\perp I}$ as a Coxeter group and a description of Y_I as the fundamental group of a 2-cell complex $\mathcal{Y}$. On the other hand, $N_W(W_I)$ is the semidirect product of W_I and a vertex group of G , the latter being described in terms of a certain complex corresponding to $\mathcal{Y}$ (temporarily denoted by $\mathcal{Y}'$). Now the preceding result gives a semidirect product decomposition of the vertex group of G , whose normal part is also a Coxeter group. However, unlike the case of $Z_W(W_I)$ where the normal part $W^{\perp I}$ is a reflection subgroup of W , the normal part for the case of $N_W(W_I)$ is in general *not* a reflection subgroup of W . The property that our normal part $W^{\perp I}$ is a reflection subgroup plays a significant role in the proof of the above-mentioned result on the action of Y_I on finite irreducible components of $W^{\perp I}$. The properties of the decomposition $W^{\perp I} \rtimes Y_I$ are not immediately inherited from those of the vertex group of G (see Remark 4.7 for details). Moreover, the structure of our complex $\mathcal{Y}$ is somewhat

simpler than their complex $\mathcal{Y}'$; namely, $\mathcal{Y}$ has no 2-cells with non-simple boundaries, while $\mathcal{Y}'$ may have such a non-simple cell. Due to these differences, our description of the structure of $Z_W(W_I)$ is still hard to derive even if that of $N_W(W_I)$ is clear.

On the other hand, for the centralizer $Z_W(W_I)$, a special case where I consists of a single generator was studied and a similar decomposition was given by Brink [4], although explicit generators of the Coxeter group part ($W^{\perp I}$ in our notation) were not given. Patrick Bahls and Mike Mihalik gave a complete description of $Z_W(W_I)$ if W is an “even” Coxeter group, namely if the product of any two distinct generators has either an even or infinite order, as is found in their preprint and a recent book by Bahls [1]. Their approach is highly different from ours. The normalizer $N_W(W_I)$ for finite W_I and its subgroups (including $Z_W(W_I)$) was also studied by Richard E. Borcherds in [3]. The commensurators of W_I in W and the centralizers of parabolic subgroups A_X in Artin-Tits groups A of certain types were examined by Luis Paris [25, 24]. Eddy Godelle [12, 13] described the normalizers, centralizers and related objects of A_X in A , and revealed some new relationship between Artin-Tits groups and Coxeter groups.

The results of this paper are applied to the author’s recent studies on the isomorphism problem in Coxeter groups [18] and some relevant topics in structural properties of Coxeter groups [20, 21]. Moreover, recently Ivonne J. Ortiz [23] used the result of (a preliminary version of) this paper for explicitly computing the lower algebraic K -theory of $\Gamma_3 = O^+(3, 1) \cap GL(4, \mathbb{Z})$ (see also [22]).

1.3 Organization of this paper

The paper is organized as follows. Section 2 is devoted to some preliminaries, and Section 3 summarizes the properties of the groupoid C that are obtained by lifting the corresponding properties of the groupoid G . Section 4 investigates the groupoid C further, enhancing its similarity with Coxeter groups noticed by Brink and Howlett in [5], and derive the decomposition of the vertex group $C_I = W^{\perp I} \rtimes Y_I$ as well as the presentations of $W^{\perp I}$ and Y_I . (Although this looks similar to the decomposition $G_I = \tilde{N}_I \rtimes M_I$ given in [5], there seems no overall relation between the structures of $\tilde{N}_I$ and $W^{\perp I}$ or between those of M_I and Y_I ; see Remark 4.7 for details.) Section 5 completes the description of the entire $Z_W(W_I)$, and gives a description of $N_W(W_I)$ based on the same argument. Section 6 carries out an explicit computation of the presentation of $Z_W(W_I)$ with an example. Finally, by virtue of the results in the preceding sections, Section 7 reveals a remarkable property (under a certain assumption on I) that the conjugation action of the subgroup Y_I on each finite irreducible component of the Coxeter group $W^{\perp I}$ becomes trivial.

Acknowledgments. This paper is an enhancement of the material in Master’s thesis [19] by the author. The author would like to express his deep gratitude to everyone who helped him, especially to Professor Itaru Terada who was the supervisor of the author during the graduate course, and to Professor Kazuhiko Koike, for their invaluable advice and encouragement. The present work was inspired by Brink’s paper [4] in the first place, and the paper [5] of Brink and Howlett incited various improvements. A part of this work was supported by JSPS Research Fellowship (No. 16-10825).

2 Preliminaries

2.1 Groupoids and related objects

We briefly summarize the notion of groupoids, fixing some notation and convention. We refer to [7] for the terminology and facts not mentioned here.

A *groupoid* is a small category whose morphisms are all invertible; namely a family of sets $G = \{G_{x,y}\}_{x,y \in I}$ with index set $\mathcal{V}(G) = I$ endowed with (1) multiplications $G_{x,y} \times G_{y,z} \rightarrow G_{x,z}$ satisfying the associativity law, (2) an identity element $1 = 1_x$ in each $G_{x,x}$, and (3) an inverse $g^{-1} \in G_{y,x}$ for every $g \in G_{x,y}$. In particular, each $G_{x,x}$ is a group, called a *vertex group* of G . We write $g \in G$ to signify $g \in \bigsqcup_{x,y \in I} G_{x,y}$, and $X \subseteq G$ for $X \subseteq \bigsqcup_{x,y \in I} G_{x,y}$. If $g \in G_{y,x}$ we call y and x the *target* and *source* of g , respectively. (Note that this is reverse to the usual convention. The reason is that we will consider later an action of an element $g \in G_{y,x}$ of a certain groupoid G from the left that maps an element associated to x to one associated to y .) An element $g \in G$ is called a *loop* in G if the source and target of g are equal, namely if g lies in some vertex group of G . A sequence $(g_l, g_{l-1}, \dots, g_1)$ of elements of G is *composable* if the source of g_{i+1} is equal to the target of g_i for every $1 \leq i \leq l-1$. A *homomorphism* between groupoids is a covariant functor between them regarded as categories. Notions such as *isomorphisms* and *subgroupoids* are defined as usual.

Let $X \subseteq G$. A *word* in the set $X^{\pm 1} = X \sqcup \{x^{-1} \mid x \in X\}$ is a composable sequence $(x_l^{\varepsilon_l}, \dots, x_2^{\varepsilon_2}, x_1^{\varepsilon_1})$ of symbols in $X^{\pm 1}$. We say that this word *represents* the element $x_l^{\varepsilon_l} \cdots x_2^{\varepsilon_2} x_1^{\varepsilon_1}$ of G . The set X is called a *generating set* of G if every element of G is represented by a word in $X^{\pm 1}$. Given a generating set X of G , a set R of formal expressions of the form $\xi = \eta$, where ξ and η are words in $X^{\pm 1}$, is called a set of *fundamental relations* of G with respect to X if (1) the two words ξ and η in each expression $\xi = \eta$ in R represent the same element of G , and (2) if two words ζ and ω in $X^{\pm 1}$ represent the same element of G , then ω is obtained from ζ by a finite steps of transformations replacing a subword matching one side of an expression in $R \cup \{x^\varepsilon x^{-\varepsilon} = \emptyset \mid x \in X \text{ and } \varepsilon \in \{\pm 1\}\}$ (where $\emptyset$ represents the empty word) by the other side (or vice versa).

Let $\mathcal{G}$ be an unoriented graph on vertex set $\mathcal{V}$ with no loops. The *fundamental groupoid* of $\mathcal{G}$, denoted here by $\pi_1(\mathcal{G}; *, *)$, is a groupoid with index set $\mathcal{V}$. The edges of $\mathcal{G}$, each endowed with one of the two possible orientations, say, from $x \in \mathcal{V}$ to $y \in \mathcal{V}$, define elements of $\pi_1(\mathcal{G}; y, x) = \pi_1(\mathcal{G}; *, *)_{y,x}$, and $\pi_1(\mathcal{G}; *, *)$ is freely generated by these elements. Note that the same edge endowed with two opposite orientations give inverse elements to each other. A vertex group $\pi_1(\mathcal{G}; x) = \pi_1(\mathcal{G}; x, x)$ is the *fundamental group* of $\mathcal{G}$ at x , which is a free group. To keep consistency with our convention for groupoids, in this paper we write a path in a graph from right to left, hence a path $e_l \cdots e_2 e_1$, where e_i is an oriented edge from x_{i-1} to x_i for $1 \leq i \leq l$, determines an element of $\pi_1(\mathcal{G}; x_l, x_0)$.

In this paper, the term “*complex*” refers to a pair $\mathcal{K} = (\mathcal{K}^1, \mathcal{C})$ of a graph $\mathcal{K}^1$, called the 1-skeleton of $\mathcal{K}$, and a set $\mathcal{C}$ of some (not necessarily simple) closed paths in $\mathcal{K}$. As an analogy of usual 2-dimensional cell complexes, each closed path $c \in \mathcal{C}$ is regarded as a “2-cell” of $\mathcal{K}$ with boundary c . When $\mathcal{K}^1$ has no loops, the *fundamental groupoid* of $\mathcal{K}$, denoted similarly by $\pi_1(\mathcal{K}; *, *)$, is constructed from $\pi_1(\mathcal{K}^1; *, *)$ by adding new relations “boundary of a 2-cell = 1”. Note that to every complex $\mathcal{K}$ a topological space (called a *geometric realization* of $\mathcal{K}$) is associated in such a way that $\pi_1(\mathcal{K}; *, *)$ is naturally a full subgroupoid of the fundamental groupoid of the space (see [7, Section 5.6] for details).

A groupoid $\tilde{G}$ is called a *covering groupoid* of a groupoid G with *covering map* $p : \tilde{G} \rightarrow G$ if p is a surjective homomorphism and, moreover, for any $g \in G$ with source x and $\tilde{x} \in \mathcal{V}(\tilde{G})$ projecting onto x (called a *lift* of x), there exists a unique element $\tilde{g} \in \tilde{G}$ with source $\tilde{x}$ projecting onto g (also called a *lift* of g); see e.g., [6, Section 9.2]. Any lift of an identity element in G is an identity element in $\tilde{G}$. If $(g_l, \dots, g_2, g_1)$ is a composable sequence of elements of G , then for any lift $\tilde{x}$ of the source x of g_1 , there is a unique composable sequence $(\tilde{g}_l, \dots, \tilde{g}_2, \tilde{g}_1)$ of lifts of $g_l, \dots, g_2, g_1$ respectively, with $\tilde{x}$ being the source of $\tilde{g}_1$; and the product $\tilde{g}_l \cdots \tilde{g}_2 \tilde{g}_1$ is the lift of $g_l \cdots g_2 g_1$ with source $\tilde{x}$. In later sections we will use the following property of covering groupoids, whose proof is straightforward and so omitted here:

Proposition 2.1. *Suppose that a groupoid G admits a covering groupoid $\tilde{G}$, a generating set X and a set R of fundamental relations. Then the set $\tilde{X}$ of the lifts of the generators $g \in X$ generates $\tilde{G}$. Moreover, $\tilde{G}$ admits, with respect to the $\tilde{X}$, a set $\tilde{R}$ of fundamental relations consisting of the formal expressions $\tilde{\xi} = \tilde{\eta}$ such that $\tilde{\xi}$ and $\tilde{\eta}$ are words in $\tilde{X}^{\pm 1}$ with the same source and these are lifts of corresponding terms of an expression $\xi = \eta$ belonging to R .*

2.2 Coxeter groups

The basics of Coxeter groups summarized here are found in [16] unless otherwise noticed.

2.2.1 Definitions

A pair (W, S) of a group W and its generating set S is called a *Coxeter system* if W admits the following presentation

$$W = \langle S \mid (st)^{m_{s,t}} = 1 \text{ for all } s, t \in S \text{ such that } m_{s,t} < \infty \rangle ,$$

where the $m_{s,t} \in \{1, 2, \dots\} \cup \{\infty\}$ are symmetric in s and t , and $m_{s,t} = 1$ if and only if $s = t$. Note that *the set S may have infinite cardinality*. A group W is called a *Coxeter group* if (W, S) is a Coxeter system for some $S \subseteq W$. An *isomorphism of Coxeter systems* from (W, S) to (W', S') is a group isomorphism $W \rightarrow W'$ that maps S onto S' . It is shown that $m_{s,t}$ is precisely the order of $st \in W$, therefore the system (W, S) determines uniquely the data $(m_{s,t})_{s,t}$ and hence the *Coxeter graph* Γ , which is a simple unoriented graph with vertex set S in which two vertices $s, t \in S$ are joined by an edge with label $m_{s,t}$ if and only if $m_{s,t} \geq 3$ (by usual convention, the label is omitted when $m_{s,t} = 3$). For $I \subseteq S$, the subgroup $W_I = \langle I \rangle$ of W generated by I is called a *parabolic subgroup*. If I is (the vertex set of) a connected component of Γ , then W_I and I are called an *irreducible component* of W (or of (W, S) , if we emphasize the generating set S) and of S , respectively. Now W is the (restricted) direct product of its irreducible components. If Γ is connected, then (W, S) , W and S are called *irreducible*. If $I \subseteq S$, then (W_I, I) is also a Coxeter system, of which the Coxeter graph Γ_I is the full subgraph of Γ with vertex set I and the length function is the restriction of the length function ℓ of (W, S) .

2.2.2 Geometric representation and root systems

Let V denote the space of the *geometric representation* of (W, S) over $\mathbb{R}$ equipped with a basis $\Pi = \{\alpha_s \mid s \in S\}$ and a W -invariant symmetric bilinear form $\langle \cdot, \cdot \rangle$ determined by

$$\langle \alpha_s, \alpha_t \rangle = \begin{cases} -\cos(\pi/m_{s,t}) & \text{if } m_{s,t} < \infty ; \\ -1 & \text{if } m_{s,t} = \infty , \end{cases}$$

where W acts faithfully on V by $s \cdot v = v - 2\langle \alpha_s, v \rangle \alpha_s$ for $s \in S$ and $v \in V$. Then the *root system* $\Phi = W \cdot \Pi$ consists of unit vectors with respect to the $\langle \cdot, \cdot \rangle$, and it is the disjoint union $\Phi = \Phi^+ \sqcup \Phi^-$ of $\Phi^+ = \Phi \cap \mathbb{R}_{\geq 0} \Pi$ and $\Phi^- = -\Phi^+$ where $\mathbb{R}_{\geq 0} \Pi$ signifies the set of nonnegative linear combinations of elements of Π . Elements of Φ , Φ^+ , and Φ^- are called *roots*, *positive roots*, and *negative roots*, respectively. For any subset $\Psi \subseteq \Phi$ and $w \in W$, put

$$\Psi^\pm = \Psi \cap \Phi^\pm , \text{ respectively,}$$

and

$$\Psi[w] = \{\gamma \in \Psi^+ \mid w \cdot \gamma \in \Phi^-\} .$$

We have $\ell(w) = |\Phi[w]|$, hence $w = 1$ if and only if $\Phi[w] = \emptyset$. This implies that the set $\Phi[w]$ determines the element $w \in W$ uniquely. The following property is well known:

Lemma 2.2. For $w_1, w_2 \in W$, the following conditions are equivalent:

1. $\ell(w_1 w_2) = \ell(w_1) + \ell(w_2)$;
2. $\Phi[w_2] \subseteq \Phi[w_1 w_2]$;
3. $\Phi[w_1] \cap \Phi[w_2^{-1}] = \emptyset$;
4. $\Phi[w_1 w_2] = (w_2^{-1} \cdot \Phi[w_1]) \sqcup \Phi[w_2]$.

For any element $v = \sum_{s \in S} c_s \alpha_s \in V$, define the *support* of v by

$$\text{Supp } v = \{s \in S \mid c_s \neq 0\} \text{ .}$$

For each $I \subseteq S$, put

$$\Pi_I = \{\alpha_s \mid s \in I\} \subseteq \Pi, \quad V_I = \text{Supp } \Pi_I \subseteq V \text{ and } \Phi_I = \Phi \cap V_I \text{ .}$$

It is well known that Φ_I coincides with the root system $W_I \cdot \Pi_I$ of (W_I, I) (see also Theorem 2.3(3) below). The support of any $\gamma \in \Phi$ is irreducible; this follows from the facts that we have $\gamma \in \Phi_I$ where $I = \text{Supp } \gamma$, and that $\text{Supp}(w \cdot \alpha_s)$ (for $w \in W_I$ and $s \in I$) is contained in the irreducible component of I containing s .

2.2.3 Reflection subgroups

For a $\gamma = w \cdot \alpha_s \in \Phi$, let $s_\gamma = w s w^{-1}$ denote the *reflection* along γ acting on V by $s_\gamma \cdot v = v - 2\langle \gamma, v \rangle \gamma$ for $v \in V$. Note that $\ell(us_\gamma) \neq \ell(u)$ for any $u \in W$. For any subset $\Psi \subseteq \Phi$, let $W(\Psi)$ denote the subgroup generated by all s_γ with $\gamma \in \Psi$. A subgroup of W of this form is called a *reflection subgroup*. Note that $W(\Psi) = W(W(\Psi) \cdot \Psi)$ and $-W(\Psi) \cdot \Psi = W(\Psi) \cdot \Psi$. Moreover, let $\Pi(\Psi)$ denote the set of all “simple roots” $\gamma \in (W(\Psi) \cdot \Psi)^+$, namely all the γ such that any expression $\gamma = \sum_{i=1}^r c_i \beta_i$ with $c_i > 0$ and $\beta_i \in (W(\Psi) \cdot \Psi)^+$ satisfies that $\beta_i = \gamma$ for all i . Put

$$S(\Psi) = \{s_\gamma \mid \gamma \in \Pi(\Psi)\} \text{ .}$$

It was shown by Vinay V. Deodhar [10] that $(W(\Psi), S(\Psi))$ is a Coxeter system. Note that Matthew Dyer [11] also proved independently that $W(\Psi)$ is a Coxeter group and determined a corresponding generating set, which in fact coincides with $S(\Psi)$. First, we summarize Deodhar’s result (the claim 1 below) and some related properties:

Theorem 2.3. Let $\Psi \subseteq \Phi$ be any subset.

1. The pair $(W(\Psi), S(\Psi))$ is a Coxeter system. For any $\beta \in \Pi(\Psi)$, the reflection s_β permutes the elements of $(W(\Psi) \cdot \Psi)^+ \setminus \{\beta\}$. Moreover, any $\gamma \in (W(\Psi) \cdot \Psi)^+$ can be expressed as a nonnegative linear combination of elements of $\Pi(\Psi)$.
2. Let $\beta, \gamma \in W(\Psi) \cdot \Psi$ and $w \in W(\Psi)$. Then $ws_\beta w^{-1} = s_\gamma$ if and only if $w \cdot \beta = \pm \gamma$.
3. If $\gamma \in \Phi$ and $s_\gamma \in W(\Psi)$, then $\gamma \in W(\Psi) \cdot \Pi(\Psi)$. In particular, we have $W(\Psi) \cdot \Psi = W(\Psi) \cdot \Pi(\Psi)$.
4. Let ℓ_Ψ be the length function of $(W(\Psi), S(\Psi))$. Then for $w \in W(\Psi)$ and $\gamma \in (W(\Psi) \cdot \Psi)^+$, we have $\ell_\Psi(ws_\gamma) < \ell_\Psi(w)$ if and only if $w \cdot \gamma \in \Phi^-$.
5. If $w \in W(\Psi)$, then $\ell_\Psi(w) = |(W(\Psi) \cdot \Psi)[w]| < \infty$.

Note that the claim 2 is well known for arbitrary roots. On the other hand, once the claim 3 is proven, the claims 4 and 5 will follow from the proof of the main theorem of Deodhar's another paper [9]. To prove the claim 3, we require another theorem of Deodhar [8]:

Theorem 2.4 ([8, Theorem 5.4]). *If $w \in W$ and $w^2 = 1$, then w decomposes into a product of commuting reflections along pairwise orthogonal roots. Hence if w is an involution, then $w \cdot \gamma = -\gamma$ for some $\gamma \in \Phi$.*

Proof of Theorem 2.3(3). For the first part, let $\gamma \in \Phi$ such that $s_\gamma \in W(\Psi)$. Then Theorem 2.4 gives us a decomposition $s_\gamma = s_{\beta_1} \cdots s_{\beta_r}$ of the involution s_γ in a Coxeter group $W(\Psi)$ into pairwise commuting distinct reflections s_{β_i} along $\beta_i \in W(\Psi) \cdot \Pi(\Psi)$. Write $\gamma = w \cdot \alpha_s$ with $s \in S$ and $w \in W$, then $s = w^{-1} s_\gamma w = s_{\beta'_1} \cdots s_{\beta'_r}$ where all $\beta'_i = w^{-1} \cdot \beta_i$ are orthogonal as well as the β_i . This implies that $s \cdot \beta'_i = -\beta'_i$, therefore $\beta'_i = \pm \alpha_s$ for every i . Hence we have $r = 1$ and $\gamma = \pm w \cdot \beta'_1 = \pm \beta_1 \in W(\Psi) \cdot \Pi(\Psi)$, as desired.

For the second part, any element $\gamma \in W(\Psi) \cdot \Psi$ satisfies that $s_\gamma \in W(W(\Psi) \cdot \Psi) = W(\Psi)$, therefore $\gamma \in W(\Psi) \cdot \Pi(\Psi)$ by the first part. The other inclusion is obvious. $\square$

Secondly, we summarize a part of Dyer's result and its consequences needed later. We say that a subset $\Psi \subseteq \Phi^+$ is a *root basis* if for all $\beta, \gamma \in \Psi$, we have

$$\begin{cases} \langle \beta, \gamma \rangle = -\cos(\pi/m) & \text{if } s_\beta s_\gamma \text{ has order } m < \infty ; \\ \langle \beta, \gamma \rangle \leq -1 & \text{if } s_\beta s_\gamma \text{ has infinite order.} \end{cases}$$

Theorem 2.5 ([11, Theorem 4.4]). *Let $\Psi \subseteq \Phi^+$ be any subset. Then we have $\Pi(\Psi) = \Psi$ if and only if Ψ is a root basis.*

Corollary 2.6. *Let $\Psi \subseteq \Phi^+$ be a root basis such that $|W(\Psi)| < \infty$. Then Ψ is a basis of a positive definite subspace of V with respect to the form $\langle \cdot, \cdot \rangle$.*

Proof. By Theorem 2.5, the finite Coxeter group $W(\Psi)$ has its own geometric representation space V' with basis $\Pi' = \{\bar{\beta} \mid \beta \in \Psi\}$ and positive definite bilinear form $\langle \cdot, \cdot \rangle'$ (see [16, Theorem 6.4]) such that $\langle \beta, \gamma \rangle = \langle \bar{\beta}, \bar{\gamma} \rangle'$ for $\beta, \gamma \in \Psi$. Now if $v = \sum_{\beta \in \Psi} c_\beta \beta \in \text{span } \Psi$ and $\langle v, v \rangle \leq 0$, then $\langle \bar{v}, \bar{v} \rangle' = \langle v, v \rangle \leq 0$ where $\bar{v} = \sum_{\beta \in \Psi} c_\beta \bar{\beta} \in V'$, implying that $c_\beta = 0$ as desired. $\square$

Here we present two more results needed later as well. The first one is a slight improvement of [15, Proposition 2.6]. Note that the assumption $|S| < \infty$ in the original statement is in fact not necessary; namely we have:

Proposition 2.7. *Let $\Psi \subseteq \Phi^+$ be a root basis such that $|W(\Psi)| < \infty$, and $U = \text{span } \Psi$. Then there exist $w \in W$ and $I \subseteq S$ such that $|W_I| < \infty$ and $w \cdot (U \cap \Phi^+) = \Phi_I^+$. Moreover, this w maps $U \cap \Pi$ into Π_I .*

Proof. First, Theorem 2.3(3) implies that the set $\{\gamma \in \Phi \mid s_\gamma \in W(\Psi)\}$ coincides with the root system $W(\Psi) \cdot \Psi$ of $W(\Psi)$, which is finite by the hypothesis. Thus the hypothesis of the original proposition [15, Proposition 2.6] is satisfied, hence it follows from that proposition that there are $w \in W$ and $I \subseteq S$ such that $|W_I| < \infty$ and $w \cdot (U \cap \Phi) = \Phi_I$ (note that $U = \text{span}(W(\Psi) \cdot \Psi)$). Choosing such a w with shortest length, it also holds that $w^{-1} \cdot \Phi_I^+ \subseteq U \cap \Phi^+$. Indeed, if $\gamma \in \Phi_I[w^{-1}]$, then $s_\gamma w \in W$ also satisfies the condition and is shorter than w (apply Theorem 2.3(4) to w^{-1}). Hence the first claim holds.

For the second claim, if $\alpha_s \in U \cap \Pi$ and $w \cdot \alpha_s = \sum_{t \in I} c_t \alpha_t$ with $c_t \geq 0$, then $\alpha_s = \sum_{t \in I} c_t w^{-1} \cdot \alpha_t$, while $w^{-1} \cdot \alpha_t \in \Phi^+$. Thus we have $w^{-1} \cdot \alpha_t = \alpha_s$ whenever $c_t > 0$, showing that $w \cdot \alpha_s = \alpha_t \in \Pi_I$ as desired. $\square$

The second result is that the theorem [16, Theorem 1.12 (d)] also holds for a general W :

Proposition 2.8. *Let $\Psi \subseteq \Phi$ be any subset. Then any $w \in W$ of finite order that fixes Ψ pointwise decomposes into a product of reflections also having this property.*

Proof. First, a well-known theorem of Jacques Tits implies that the finite subgroup $\langle w \rangle$ of W is conjugate to a subgroup of a finite W_I (see [2, Theorem 4.5.3] for a proof), therefore we may assume without loss of generality that $w \in W_I$. Since V_I is positive definite (see [16, Theorem 6.4]), each $\gamma \in \Psi$ decomposes as $\gamma = v + v'$ with $v \in V_I$ and $v' \in V_I^\perp$. Now $w \in W_I$ fixes every $v = \gamma - v'$, therefore the result for the finite W_I yields a decomposition $w = s_{\beta_1} \cdots s_{\beta_r}$, where each reflection $s_{\beta_j} \in W_I$ fixes every v , hence fixes every $\gamma = v + v' \in \Psi$, as desired. $\square$

2.2.4 Finite parabolic subgroups and their longest elements

We say that a subset $I \subseteq S$ is of *finite type* if $|W_I| < \infty$, or equivalently $|\Phi_I| < \infty$. The finite irreducible Coxeter groups have been classified as summarized in [16, Chapter 2]. Here we give a canonical labelling $r_1, r_2, \dots, r_n$ (where $n = |I|$) of elements of an irreducible subset $I \subseteq S$ of each finite type used in this paper in the following manner, where we put $m_{i,j} = m_{r_i, r_j}$ for simplicity and the $m_{i,j}$ not listed here are all equal to 2:

Type A_n ($n \geq 2$): $m_{i,i+1} = 3$ ($1 \leq i \leq n-1$);

Type B_n ($n \geq 2$): $m_{i,i+1} = 3$ ($1 \leq i \leq n-2$) and $m_{n-1,n} = 4$;

Type D_n ($n \geq 4$): $m_{i,i+1} = m_{n-2,n} = 3$ ($1 \leq i \leq n-2$);

Type E_n ($n = 6, 7, 8$): $m_{1,3} = m_{2,4} = m_{i,i+1} = 3$ ($3 \leq i \leq n-1$);

Type F_4 : $m_{1,2} = m_{3,4} = 3$ and $m_{2,3} = 4$;

Type H_n ($n = 3, 4$): $m_{1,2} = 5$ and $m_{i,i+1} = 3$ ($2 \leq i \leq n-1$);

Type $I_2(m)$ ($m \geq 5$): $m_{1,2} = m$.

Let $w_0(I)$ denote the (unique) longest element of a finite W_I , which has order two and maps Π_I onto $-\Pi_I$. Now let I be irreducible of finite type. If I is of type A_n ($n \geq 2$), D_k (k odd), E_6 or $I_2(m)$ (m odd), then the automorphism of the Coxeter graph Γ_I of W_I induced by (the conjugation action of) $w_0(I)$ is the unique nontrivial automorphism on Γ_I . Otherwise $w_0(I)$ lies in the center $Z(W_I)$ of W_I and the induced automorphism on Γ_I is trivial, in which case we say that I is of (-1) -type. Moreover, if W_I is finite but not irreducible, then $w_0(I) = w_0(I_1) \cdots w_0(I_k)$ where the I_i are the irreducible components of I . Note that for an arbitrary $I \subseteq S$, the center $Z(W_I)$ is an elementary abelian 2-group generated by the $w_0(J)$ where J runs over all irreducible components of I of (-1) -type.

3 The groupoid C

From now on, we fix an arbitrarily given subset $I \subseteq S$ unless specifically noted otherwise. In this section, we introduce and study a groupoid C , one of whose vertex groups gives a fairly large part (still not all) of the centralizer $Z_W(W_I)$. As we shall see in Section 3.3, our groupoid C is a covering of the groupoid G defined by Brink and Howlett in [5] for their study of the normalizer $N_W(W_I)$. As a consequence, many properties, but not all, of C can be obtained from those of G .

Here we survey the connection among the centralizer $Z_W(W_I)$ and the two groupoids C and G . If X is a W -set (that is, a set with W -action) and $Y \subseteq X$, let $\mathcal{G} = (X; Y)$ denote tentatively a groupoid with index set Y defined by $\mathcal{G} = \{\mathcal{G}_{y,y'}\}_{y,y' \in Y}$ and $\mathcal{G}_{y,y'} = \{w \in W \mid y = w \cdot y'\}$, with multiplication induced by that of W . The vertex group $\mathcal{G}_{y,y}$ of $\mathcal{G}$ at $y \in Y$ is the stabilizer of y in W . If X' is another W -set and $Y' \subseteq X'$, then a W -equivalent map $\varphi : X \rightarrow X'$ with $\varphi(Y) \subseteq Y'$ induces a groupoid homomorphism $\varphi_* : (X; Y) \rightarrow (X'; Y')$. It is a covering of groupoids if $\varphi(Y) = Y'$ and $Y = \varphi^{-1}(Y')$.

Now fix a set Λ with $|\Lambda| = |I|$. For a W -set X , let X^* temporarily denote the set of all injections $\Lambda \rightarrow X$ with the W -action induced from that on X . Then the centralizer $Z_W(W_I)$, that is the final target of our study, is the vertex group of a groupoid $(T^*; S^*)$ at an element $x_I \in S^*$, where T denotes the set of reflections in W with respect to S (on which W acts by conjugation) and the image of the map x_I is I .

Look at the following diagram of groupoids, whose morphisms are induced by the obvious W -equivalent maps. Here the symbol $\binom{X}{|\Lambda|}$, where X is a W -set, denotes the set of all subsets of X with cardinality $|\Lambda|$ with the W -action induced from that on X . The two morphisms marked $*$ are coverings, and the one marked $\#$ is an inclusion as a full subgroupoid. Note that the composition of the vertical arrows induces a bijection on the index sets. The groupoid C we introduce below is isomorphic to the connected component of the groupoid $(\Phi^*; \Pi^*)$ on the top-left corner containing the lift of x_I . Since $Z_W(W_I)$ contains elements projected from those of the groupoid $(\Phi^*; (\pm\Pi)^*)$ with source being the lift of $x_I \in S^*$ that lies in $\Pi^* \subseteq (\pm\Pi)^*$ and target being its other lifts in $(\pm\Pi)^* \setminus \Pi^*$, the vertex group C_I is only a part of $Z_W(W_I)$ in general. On the other hand, the groupoid G in [5] is a connected component of the groupoid $((\Phi_\Lambda); (\Pi_\Lambda))$ on the right.

$$\begin{array}{ccccc}
C_I & \subseteq & (\Phi^*; \Pi^*) & \xrightarrow{*} & ((\Phi_\Lambda); (\Pi_\Lambda)) \supseteq G \\
& & \# \downarrow & & \\
& & (\Phi^*; (\pm\Pi)^*) & & \\
& & * \downarrow & & \\
Z_W(W_I) & \subseteq & (T^*; S^*) & &
\end{array}$$

3.1 Definitions

In what follows, we would like to deal with an “ordered tuple” consisting of the elements of a given subset of S . The reason is that elements of the centralizer $Z_W(W_I)$ not only leaves the set I invariant but also fix every element of I pointwise. For the purpose, first we fix an auxiliary index set Λ_0 having the same cardinality as S . Then put

$$S^{(\Lambda)} = \{x : \Lambda \rightarrow S \mid x \text{ is injective}\} \text{ for each } \Lambda \subseteq \Lambda_0$$

and

$$S^* = \bigcup_{\Lambda \subseteq \Lambda_0} S^{(\Lambda)} .$$

For $x \in S^{(\Lambda)}$ and $\lambda \in \Lambda$, we write x_λ for $x(\lambda)$; thus x may be regarded as a duplicate-free “ Λ -tuple” $(x_\lambda) = (x_\lambda)_{\lambda \in \Lambda}$ of elements of S . If $|\Lambda| = n < \infty$, then it is regarded as a duplicate-free sequence $(x_1, x_2, \dots, x_n)$ by identifying Λ with $\{1, 2, \dots, n\}$. We write

$$x_A = \{x_\lambda \mid \lambda \in A\} \text{ for each } A \subseteq \Lambda, \text{ and } [x] = x_\Lambda .$$

If we put

$$\widehat{C}_{x,y} = \{w \in W \mid \alpha_{x_\lambda} = w \cdot \alpha_{y_\lambda} \text{ for all } \lambda \in \Lambda\} \text{ for all } x, y \in S^{(\Lambda)}$$

and $\widehat{C}_{x,y} = \emptyset$ for all $x \in S^{(\Lambda)}$ and $y \in S^{(\Lambda')}$ such that $\Lambda \neq \Lambda'$, then $\widehat{C} = \{\widehat{C}_{x,y}\}_{x,y \in S^*}$ is a groupoid with vertex set $\mathcal{V}(\widehat{C}) = S^*$ and multiplication induced by that of W . Write

$$x = w \cdot y \text{ if } x, y \in S^* \text{ and } w \in \widehat{C}_{x,y} . \quad (3.1)$$

Now we fix elements

$$\Lambda \subseteq \Lambda_0 \text{ and } x_I \in S^{(\Lambda)} \text{ such that } [x_I] = I ,$$

and define the groupoid C as the connected component of $\widehat{C}$ containing x_I ; hence

$$\mathcal{V}(C) = \{x \in S^{(\Lambda)} \mid \widehat{C}_{x,x_I} \neq \emptyset\} \text{ and } C_{x,y} = \widehat{C}_{x,y} \text{ for } x, y \in \mathcal{V}(C) .$$

Note that $C_{x,y} \neq \emptyset$ for all $x, y \in \mathcal{V}(C)$. We put

$$C_I = C_{x_I, x_I} = \{w \in W \mid w \cdot \alpha_s = \alpha_s \text{ for all } s \in I\} ,$$

which is a normal subgroup of $Z_W(W_I)$. We will see below that this C_I occupies a fairly large part, but in general not the whole, of $Z_W(W_I)$.

We will see in Theorem 4.6 the following semidirect product decomposition

$$C_I = W^{\perp I} \rtimes Y_I . \quad (3.2)$$

We define the factors below. First, for arbitrary subsets $I, J \subseteq S$, let

$$\Phi_J^{\perp I} = \{\gamma \in \Phi_J \mid \gamma \text{ is orthogonal to every } \alpha_s \in \Pi_I\}$$

and

$$W_J^{\perp I} = W(\Phi_J^{\perp I})$$

(see Section 2.2.3 for notations). Note that for any $\gamma \in \Phi_J^{\perp I}$, we have $s_\gamma \cdot \alpha_s = \alpha_s$ for every $s \in I$, therefore $s_\gamma \cdot \Phi_J^{\perp I} = \Phi_J^{\perp I}$. Hence by Theorem 2.3, $W_J^{\perp I}$ is a Coxeter group with root system $W_J^{\perp I} \cdot \Phi_J^{\perp I} = \Phi_J^{\perp I}$, generating set and simple system given by

$$R^{J,I} = S(\Phi_J^{\perp I}) \text{ and } \Pi^{J,I} = \Pi(\Phi_J^{\perp I}) , \text{ respectively.}$$

In the notations, the symbol J will be omitted when $J = S$. In particular, the factor $W^{\perp I}$ in (3.2) is given by

$$W^{\perp I} = W_S^{\perp I} = \langle \{s_\gamma \mid \gamma \in \Phi^{\perp I}\} \rangle .$$

For the factor Y_I in (3.2), first define a subgroupoid $\widehat{Y} = \{\widehat{Y}_{x,y}\}_{x,y \in S^*}$ of $\widehat{C}$ by

$$\widehat{Y}_{x,y} = \{w \in \widehat{C}_{x,y} \mid w \cdot (\Phi^{\perp[y]})^+ \subseteq \Phi^+\} = \{w \in \widehat{C}_{x,y} \mid (\Phi^{\perp[x]})^+ = w \cdot (\Phi^{\perp[y]})^+\}$$

for $x, y \in S^*$. Here the second equality is a consequence of the following fact

$$\Phi^{\perp[y]} = w \cdot \Phi^{\perp[x]} \text{ for all } w \in \widehat{C}_{x,y} ,$$

also implying that $W^{\perp[x]}$ is normal in $\widehat{C}_{x,x}$. Now let Y be the full subgroupoid of $\widehat{Y}$ with vertex set $\mathcal{V}(C)$, namely

$$\mathcal{V}(Y) = \mathcal{V}(C) \text{ and } Y_{x,y} = \widehat{Y}_{x,y} \text{ for } x, y \in \mathcal{V}(C) ,$$

and write

$$Y_I = Y_{x_I, x_I} = \{w \in C_I \mid (\Phi^{\perp I})^+ = w \cdot (\Phi^{\perp I})^+\} .$$

Then Y is a subgroupoid of C . Note that $W^{\perp[x]} \cap Y_{x,x} = 1$ since any element of $W^{\perp[x]} \cap Y_{x,x}$ has length zero by virtue of Theorem 2.3(5).

3.2 Preceding results on normalizers

Here we summarize some preceding results on the normalizers $N_W(W_I)$ of W_I in W given by Brink and Howlett [5]. Their first step is the decomposition $N_W(W_I) = W_I \rtimes N_I$ (see [5, Proposition 2.1]), where the factor

$$N_I = \{w \in W \mid w \cdot \Pi_I = \Pi_I\}$$

is the main subject of the paper [5].

The following theorem, used in [5] for general S , is proven in [8] by Deodhar under the assumption $|S| < \infty$. A proof for the general case is given in [21].

Theorem 3.1 (Deodhar). *Let $K \subseteq S$ be any subset and $J \subset K$ a proper subset, and suppose that W_K is infinite and irreducible. Then $|\Phi_K \setminus \Phi_J| = \infty$.*

For $J, K \subseteq S$, let $J_{\sim K}$ denote the union of the connected components of the graph $\Gamma_{J \cup K}$ having nonempty intersection with K . Then Theorem 3.1 implies that, in the case $J \cap K = \emptyset$, the set $J_{\sim K} \subseteq S$ is of finite type if and only if the set $\Phi_{J \cup K} \setminus \Phi_J = \Phi_{J_{\sim K}} \setminus \Phi_J$ is finite. Here the last equality follows from the irreducibility of the support of any root (see Section 2.2.2). In this case, it is shown in [5] that

$$w_0(J_{\sim K})w_0(J_{\sim K} \setminus K) \cdot \Pi_J \subseteq \Pi_{J \cup K}$$

and

$$\Phi[w_0(J_{\sim K})w_0(J_{\sim K} \setminus K)] = \Phi_{J \cup K}^+ \setminus \Phi_J .$$

Moreover, it is easy to show that

$$(w_0(J_{\sim K})w_0(J_{\sim K} \setminus K))^{-1} = w_0(J'_{\sim K'})w_0(J'_{\sim K'} \setminus K') , \quad (3.3)$$

where we put $\Pi_{J'} = w_0(J_{\sim K})w_0(J_{\sim K} \setminus K) \cdot \Pi_J$ and $K' = (J \cup K) \setminus J'$.

An element $u \in W$ is called a *right divisor* of $w \in W$ if $\ell(w) = \ell(wu^{-1}) + \ell(u)$. The following lemma is a generalization of [5, Lemma 4.1]. The proof of the original lemma can be easily adapted.

Lemma 3.2 (See [5, Lemma 4.1]). *Let $w \in W$ and $J, K \subseteq S$, and suppose that $w \cdot \Pi_J \subseteq \Pi$ and $w \cdot \Pi_K \subseteq \Phi^-$. Then $J \cap K = \emptyset$, the set $J_{\sim K}$ is of finite type and $w_0(J_{\sim K})w_0(J_{\sim K} \setminus K)$ is a right divisor of w .*

A groupoid G played a central role in the paper [5]. We recall the definition. Given $I \subseteq S$, put

$$\begin{aligned} \mathcal{J} &= \{J \subseteq S \mid \Pi_J = w \cdot \Pi_I \text{ for some } w \in W\} ; \\ G_{J,K} &= \{(J, w, K) \mid w \in W \text{ and } \Pi_J = w \cdot \Pi_K\} \text{ for } J, K \in \mathcal{J} . \end{aligned}$$

The multiplication in the groupoid $G = \{G_{J,K}\}_{J,K \in \mathcal{J}}$ is defined by

$$(J_1, w, J_2)(J_2, u, J_3) = (J_1, wu, J_3) .$$

Note that the map $(I, w, I) \mapsto w$ from $G_{I,I}$ to N_I is a group isomorphism. Now for $J \subseteq S$ and $s \in S \setminus J$, write

$$v[s, J] = w_0(J_{\sim s})w_0(J_{\sim s} \setminus \{s\}) \text{ if } J_{\sim s} \text{ is of finite type,}$$

hence $v[s, J] \cdot \Pi_J = \Pi_K$ for a unique $K \subseteq S$. Moreover, an expression $g = g_1 g_2 \cdots g_n$ of $g = (J_0, w, J_n) \in G_{J_0, J_n}$ with $g_i = (J_{i-1}, v[s_i, J_i], J_i)$ is called a *standard expression in G* if $\ell(w) = \sum_{i=1}^n \ell(v[s_i, J_i])$.

The argument in [5] requires the following theorem of Deodhar:

Theorem 3.3 ([8, Proposition 5.5]). *Any $(J, w, K) \in G$ admits a standard expression in G . Moreover, if $s \in S$ and $w \cdot \alpha_s \in \Phi^-$, then this expression can be chosen in such a way that it ends with $(J', v[s, K], K)$ for some J' .*

Now we summarize some results of [5] required in this paper:

Theorem 3.4. 1. ([5, Theorem 2.4]) *Suppose that $s, t \notin J \subseteq S$, $s \neq t$ and $J' = J_{\sim\{s,t\}}$ is of finite type, hence $w = w_0(J')w_0(J' \setminus \{s, t\})$ maps Π_J onto a unique Π_K . Then $(K, w, J) \in G$ admits exactly two standard expressions of the form*

$$(J_0, v[s_1, J_1], J_1) \cdots (J_{n-1}, v[s_n, J_n], J_n)$$

in G . Both of them consist of the same number n of factors and satisfy that $J_i \cup \{s_i\} \subseteq J \cup \{s, t\}$ for all i . Moreover, one of them satisfies $s_n = s$ and the other satisfies $s_n = t$.

2. ([5, Theorem A]) *The groupoid G is generated by the elements $(J, v[s, K], K)$. Moreover, the following two kinds of relations*

- $(J, v[s, K], K)(K, v[t, J], J) = 1$, with $K \cup \{s\} = J \cup \{t\}$,
- $g_1 g_2 \cdots g_n = g'_1 g'_2 \cdots g'_n$, where $g_1 g_2 \cdots g_n$ and $g'_1 g'_2 \cdots g'_n$ are the standard expressions of a common $w_0(J')w_0(J' \setminus \{s, t\})$ such that $s, t \notin J \subseteq S$, $s \neq t$ and $J' = J_{\sim\{s,t\}}$ is of finite type,

are fundamental relations of G with respect to these generators.

Note that $g_1 \neq g'_1$ in the statement 2, since $g_n^{-1} \cdots g_2^{-1} g_1^{-1}$ and $g'_n{}^{-1} \cdots g'_2{}^{-1} g'_1{}^{-1}$ are the standard expressions of the element $(w_0(J')w_0(J' \setminus \{s, t\}))^{-1}$ (see (3.3) and the statement 1).

In this paper, we say that a generator $(J, v[s, K], K)$ of G is a *loop generator* if $J = K$, namely if it is a loop of the groupoid G (see Section 2.1 for terminology).

3.3 Lifting from G to C

Now it is straightforward to show that C is a covering groupoid of G with covering map that sends $x \in \mathcal{V}(C)$ to $[x] \in \mathcal{J}$ and $w \in C_{x,y}$ to $([x], w, [y]) \in G_{[x],[y]}$ (see Section 2.1 for the terminology). This enables us to lift up the above results on G to its covering groupoid C , as in Theorem 3.5 below (see Proposition 2.1).

We prepare notations and terminology. For a pair $\xi = (x, s)$ of $x \in \mathcal{V}(C)$ and $s \in S \setminus [x]$ such that $[x]_{\sim s}$ is of finite type, we have $y = v[s, [x]] \cdot x \in \mathcal{V}(C)$ (see (3.1) for notation), and the generator $([y], v[s, [x]], [x])$ of G has a unique lift in $C_{y,x}$. We denote this element in $C_{y,x}$ by w_x^s or w_ξ , and write

$$\varphi(\xi) = \varphi(x, s) = (y, t) ,$$

where t is the unique element of $([x] \cup \{s\}) \setminus [y]$. Theorem 3.5(3) below implies that φ is an involutive map from the set of all such pairs (x, s) to itself. Moreover, we also use the terminology “a standard expression of w in C ” in a similar way, where the elements w_x^s play the role of the $(J, v[s, K], K)$ in G . Now we have the following results on the groupoid C by applying Proposition 2.1 as mentioned above:

Theorem 3.5. 1. *Any $w \in C$ admits a standard expression in C . Moreover, if $s \in S$ and $w \cdot \alpha_s \in \Phi^-$, then this expression can be chosen in such a way that it ends with w_x^s .*

2. Suppose that $x \in \mathcal{V}(C)$, $s, t \in S \setminus [x]$, $s \neq t$ and $J = [x]_{\sim\{s,t\}}$ is of finite type, hence $w = w_0(J)w_0(J \setminus \{s, t\})$ belongs to $C_{y,x}$ for a unique $y \in \mathcal{V}(C)$. Then w admits exactly two standard expressions of the form

$$w_{z_1}^{s_1} w_{z_2}^{s_2} \cdots w_{z_n}^{s_n}$$

in C . Both of them consist of the same number n of factors and satisfy that $[z_i] \cup \{s_i\} \subseteq [x] \cup \{s, t\}$ for all i . Moreover, one of them satisfies $s_n = s$ and the other satisfies $s_n = t$.

3. The groupoid C is generated by the elements w_x^s . Moreover, the following two kinds of relations

- $w_x^s w_y^t = 1$, with $[x] \cup \{s\} = [y] \cup \{t\}$ (or equivalently, $w_\xi w_{\varphi(\xi)} = 1$),
- $c_1 c_2 \cdots c_n = c'_1 c'_2 \cdots c'_n$, where $c_1 c_2 \cdots c_n$ and $c'_1 c'_2 \cdots c'_n$ are the standard expressions of a common $w_0(J)w_0(J \setminus \{s, t\})$ such that $x \in \mathcal{V}(C)$, $s, t \in S \setminus [x]$, $s \neq t$ and $J = [x]_{\sim\{s,t\}}$ is of finite type,

are fundamental relations of C with respect to these generators. In addition, for the relation of the second type, we have $c_1 \neq c'_1$.

We use the term “loop generator” also for C ; an element w_x^s is a loop generator of C if $w_x^s \in C_{x,x}$, or equivalently $w_x^s \cdot x = x$.

We refer to any transformation of expressions in C of the form

$$w_1 c_1 c_2 \cdots c_n w_2 \rightsquigarrow w_1 c'_1 c'_2 \cdots c'_n w_2 ,$$

where $c_1 c_2 \cdots c_n$ and $c'_1 c'_2 \cdots c'_n$ are the two expressions in the second relation in Theorem 3.5(3), as a *generalized braid move* (or a *GBM* in short). Its *loop number* is defined as a half of the total number of loop generators contained in two expressions $c_1 c_2 \cdots c_n$ and $c'_1 c'_2 \cdots c'_n$. In fact, this is equal to the number of loop generators in $c_1 c_2 \cdots c_n$, or equivalently in $c'_1 c'_2 \cdots c'_n$ (see Remark 4.5 below). Now Theorem 3.5(3) says that any two expressions in C of the same element can be converted to each other by generalized braid moves together with insertions and deletions of subwords of the form $w_\xi w_{\varphi(\xi)}$. This property will be enhanced in Proposition 4.4.

3.4 The graph $\mathcal{C}$

The set of generators w_x^s of C given in Theorem 3.5(3) can be regarded as the edge set of a (connected) graph $\mathcal{C}$ with vertex set $\mathcal{V}(C)$, where w_x^s is an edge from $x \in \mathcal{V}(C)$ to $w_x^s \cdot x \in \mathcal{V}(C)$. To regard $\mathcal{C}$ as an unoriented graph, we identify each edge w_ξ with its opposite $w_{\varphi(\xi)}$. Since we write an edge or a path in $\mathcal{C}$ from right to left by the convention mentioned in Section 2.1, the paths in $\mathcal{C}$ are the expressions of elements in C . This graph $\mathcal{C}$ will be used in our argument below.

Let $x \in \mathcal{V}(C)$ and $s, t \in S \setminus [x]$ such that $s \neq t$ and $J_{\sim(J \setminus [x])}$ is of finite type, where $J = [x] \cup \{s, t\}$. Let $\mathcal{C}(J)$ be the subgraph of $\mathcal{C}$ consisting of all vertices $y \in \mathcal{V}(C)$ with $[y] \subseteq J$ and all edges $w_y^{s'}$ with $[y] \cup \{s'\} \subseteq J$. Moreover, let $\mathcal{G}$ be the connected component of $\mathcal{C}(J)$ containing x . Then it is easy to show that, for any vertex y of $\mathcal{G}$, we have $|J \setminus [y]| = 2$, $J_{\sim(J \setminus [y])} = J_{\sim(J \setminus [x])}$, and $y_\lambda = x_\lambda$ whenever $\lambda \in \Lambda$ and $x_\lambda \notin J_{\sim(J \setminus [x])}$. (Indeed, it suffices to check the property only for adjacent vertices x, y in $\mathcal{G}$.) Thus $\mathcal{G}$ is a finite graph in which every vertex is adjacent to exactly two edges, hence it is classified into the following two types:

1. A cycle without loops. We refer to any nonempty, non-backtracking closed path p in this $\mathcal{G}$ as a *circular tour*.

2. A union of two loops and a simple (possibly empty) path joining the loops (see the left-hand side of Figure 1). We say that a closed path p in this $\mathcal{G}$ is a *shuttling tour* if it visits every vertex of $\mathcal{G}$ exactly twice (except the start point of the path) and passes each of the two loops exactly once (see the right-hand side of Figure 1).

In both cases, the path p represents an element of the finite parabolic subgroup $W_{J \sim (J \setminus [x])}$, hence the order of p as an element of W is finite.



Figure 1: The component $\mathcal{G}$ of the second type

The next lemma relates the shuttling tours with the generalized braid moves:

Lemma 3.6. *Let q be a path in $\mathcal{C}$ and $k \geq 1$. Then the followings are equivalent:*

1. $q = p^k$ as paths in $\mathcal{C}$ for a shuttling tour p of order k ;
2. $q = (c'_1 \cdots c'_r)^{-1} c_1 \cdots c_r$ as paths in $\mathcal{C}$ for a generalized braid move $c_1 \cdots c_r \rightsquigarrow c'_1 \cdots c'_r$ of loop number k .

Proof. Assuming the property 1, let p correspond to the graph $\mathcal{G}$, start with $w_y^{s'}$ and end with $(w_y^{t'})^{-1}$. Then $[y]_{\sim\{s', t'\}}$ is of finite type as mentioned above, therefore the triple (y, s', t') yields a GBM $c_1 \cdots c_r \rightsquigarrow c'_1 \cdots c'_r$ such that $(c_r, c'_r) = (w_y^{s'}, w_y^{t'})$, both $c_1 \cdots c_r$ and $c'_1 \cdots c'_r$ are non-backtracking distinct paths in $\mathcal{G}$, and $c_1 \neq c'_1$. Now the shape of $\mathcal{G}$ forces the closed path $q' = (c'_1 \cdots c'_r)^{-1} c_1 \cdots c_r$ to be a power of p , say, p^n with $n > 0$. Then this GBM has loop number n , since the path p^n contains $2n$ loops.

We show that $n = k$. Since $q' = p^n$ represents an identity element in $\mathcal{C}$, the order k of p is a divisor of n . Now if $k < n$, then $k \leq n/2$ and a path p^k representing an identity element must be a subpath of the first half $c_1 \cdots c_r$ of $q' = p^n$, contradicting the standardness of $c_1 \cdots c_r$. Thus $n = k$ as desired, hence the property 2 follows.

On the other hand, assuming the property 2, let the triple (x, s, t) correspond to the GBM as in Theorem 3.5(3). Then, since the closed path $q = (c'_1 \cdots c'_r)^{-1} c_1 \cdots c_r$ contains $2k$ loops, the same argument as the first paragraph implies that q is a certain power p^n of a shuttling tour p with $n > 0$, this p has order n , and we have $n = k$ since p^n contains $2n$ loops. Hence the property 1 follows, concluding the proof of Lemma 3.6. $\square$

4 The decomposition of $\mathcal{C}$

The aim of this section is to prove the decomposition (3.2) of the group C_I and describe the factors in detail. In the course of our argument (see Section 4.1), we develop more similarities of $\mathcal{C}$ with Coxeter groups, in addition to those inherited from those of G shown in [5] and summarized above.

4.1 Similarities of $\mathcal{C}$ with Coxeter groups

We start with the following lemma, which says that any loop generator of $\mathcal{C}$ lying in $C_{x,x}$ is a reflection along a root in $\Phi^{\perp[x]}$. This property of our groupoid $\mathcal{C}$, which the groupoid G does not possess, plays an important role in our argument below.

Lemma 4.1. For $x \in \mathcal{V}(C)$ and $s \in S \setminus [x]$, the three conditions are equivalent:

1. $[x]_{\sim_s}$ is of finite type, and $\varphi(x, s) = (x, s)$, i.e., $w_x^s \in C_{x,x}$;
2. $[x]_{\sim_s}$ is of finite type, and $\Phi^{\perp[x]}[w_x^s] \neq \emptyset$, i.e., $w_x^s \notin Y$;
3. $\Phi_{[x] \cup \{s\}}^{\perp[x]} \neq \emptyset$.

If these conditions are satisfied, we have $\Phi^{\perp[x]}[w_x^s] = (\Phi_{[x] \cup \{s\}}^{\perp[x]})^+ = \{\gamma(x, s)\}$ for a unique positive root $\gamma(x, s)$ such that $s_{\gamma(x, s)} = w_x^s$.

Proof. Put $\xi = (x, s)$. First, we deduce the property 2 from the property 1. The property 1 implies that $w_\xi = w_{\varphi(\xi)}$, while $w_{\varphi(\xi)} = w_\xi^{-1}$ (see Theorem 3.5(3)), therefore $w_\xi^2 = 1$. Thus Theorem 2.4 gives us a root γ with $w_\xi \cdot \gamma = -\gamma$, which satisfies that $\gamma \in \Phi^{\perp[x]}$ since $w_\xi \in C_{x,x}$ and the form $\langle, \rangle$ is W -invariant, proving the property 2 as desired. Moreover, the property 2 implies the property 3 since $\Phi[w_\xi] \subseteq \Phi_{[x] \cup \{s\}}$. This inclusion also implies that $\Phi^{\perp[x]}[w_\xi] \subseteq (\Phi_{[x] \cup \{s\}}^{\perp[x]})^+$.

Now we deduce the property 1 from the property 3. Let $\gamma \in (\Phi_{[x] \cup \{s\}}^{\perp[x]})^+$. Then we have $s_\gamma \in C_{x,x} \cap W_{[x] \cup \{s\}}$, therefore $\emptyset \neq \Phi[s_\gamma] \subseteq \Phi_{[x] \cup \{s\}}^+ \setminus \Phi_{[x]}$, forcing the root $s_\gamma \cdot \alpha_s$ to be negative. Hence Theorem 3.5(1) gives us a standard expression of s_γ in C ending with w_ξ . Moreover, this expression consists of only the term w_ξ , since the equality $\Phi_{[x] \cup \{s\}}^+ \setminus \Phi_{[x]} = \Phi[w_\xi]$ implies that $\Phi[s_\gamma] \subseteq \Phi[w_\xi]$ and $\ell(s_\gamma) \leq \ell(w_\xi)$. This proves that $s_\gamma = w_\xi \in C_{x,x}$. Since $\gamma \in (\Phi_{[x] \cup \{s\}}^{\perp[x]})^+$ was chosen arbitrarily, the uniqueness of $\gamma = \gamma(x, s)$ and the inclusion $(\Phi_{[x] \cup \{s\}}^{\perp[x]})^+ = \{\gamma(x, s)\} \subseteq \Phi^{\perp[x]}[w_\xi]$ follow. Moreover, it also follows that $w_\xi \cdot x = x$, therefore $\varphi(\xi) = \xi$ as desired. Hence the proof of Lemma 4.1 is concluded. $\square$

The next property is an analogy of the Exchange Condition for Coxeter groups:

Lemma 4.2. Let $w = w_{\xi_1} w_{\xi_2} \cdots w_{\xi_n} w_\zeta$ be an expression in C , where $\zeta = (x, s)$ and $\varphi(\zeta) = (y, t)$, such that the expression $w_{\xi_1} \cdots w_{\xi_n}$ is standard. Then the conditions 1–3 below are equivalent in general, and moreover, the condition 4 is also equivalent to the first three when $w_\zeta = s_\gamma$ is a loop generator with γ a positive root (see Lemma 4.1):

1. the expression $w_{\xi_1} \cdots w_{\xi_n} w_\zeta$ is not standard;
2. $w_{\xi_1} \cdots w_{\xi_n} \cdot \alpha_t \in \Phi^-$;
3. $w_{\xi_1} \cdots w_{\xi_n} \in C$ admits a standard expression ending with $w_\zeta^{-1} = w_{\varphi(\zeta)}$;
4. $w_{\xi_1} \cdots w_{\xi_n} \cdot \gamma \in \Phi^-$.

Proof. Put $u = w_{\xi_1} \cdots w_{\xi_n}$. First, the condition 2 implies the condition 3 by Theorem 3.5(1), while the condition 3 implies the condition 1 since now $\ell(uw_\zeta) = \ell(u) - \ell(w_\zeta)$. We show that the condition 1 implies the condition 2. By the hypothesis and the condition 1, we have $\ell(uw_\zeta) < \ell(u) + \ell(w_\zeta)$, hence u maps some $\gamma \in \Phi[w_\zeta^{-1}] = \Phi_{[y] \cup \{t\}}^+ \setminus \Phi_{[y]}$ to a negative root (see Lemma 2.2). Since u maps $\Pi_{[y]}$ into Π , it must map α_t to a negative root, proving the condition 2 as desired.

In the special case $w_\zeta = s_\gamma$, the condition 4 implies the condition 1, since now both u and s_γ send $\gamma \in \Phi^+$ into Φ^- and hence $\ell(us_\gamma) < \ell(u) + \ell(s_\gamma)$ by Lemma 2.2. On the other hand, the condition 3 implies the condition 4 since now we have $\gamma \in \Phi[w_\zeta^{-1}] \subseteq \Phi[u]$ by Lemma 2.2 again. Hence the proof of Lemma 4.2 is concluded. $\square$

Given $w \in C$ and its standard expression, let $\text{lp}(w)$ denote the number of the loop generators contained in this expression. It will be shown in Proposition 4.3 that $\text{lp}(w)$ is well-defined regardless of the choice of the standard expression of w . The next property is an analogy of a well-known property of Coxeter groups concerning the length of w and the set $\Phi[w]$. The proof is also analogous to the case of Coxeter groups.

Proposition 4.3. *Let $w \in C_{y,x}$. Then $\text{lp}(w)$ is equal to the cardinality of the set $\Phi^{\perp[x]}[w]$, hence is determined just by w regardless of the given standard expression. If w admits a standard expression of the form $u_0 s_{\gamma_1} u_1 s_{\gamma_2} u_2 \cdots u_{n-1} s_{\gamma_n} u_n$, where each u_i contains no loop generators and each s_{γ_i} is a loop generator with γ_i positive, then*

$$\Phi^{\perp[x]}[w] = \{\beta_i = (u_i s_{\gamma_{i+1}} u_{i+1} \cdots s_{\gamma_n} u_n)^{-1} \cdot \gamma_i \mid 1 \leq i \leq n\}.$$

Proof. Put $w_{i,j} = u_i s_{\gamma_{i+1}} u_{i+1} \cdots s_{\gamma_j} u_j$ for indices $i \leq j$, and $z_i = w_{i,n} \cdot x$. First we show that all the β_i in the statement are distinct and lie in $\Phi^{\perp[x]}[w]$, proving that $n \leq |\Phi^{\perp[x]}[w]|$. Note that $\beta_i \in \Phi^{\perp[x]}$ since $\gamma_i \in \Phi^{\perp[z_i]}$. Now if $1 \leq i < j \leq n$ and $\beta_i = \beta_j$, then we have

$$w_{i,j-1}^{-1} \cdot \gamma_i = s_{\gamma_j} w_{j,n} \cdot \beta_i = s_{\gamma_j} w_{j,n} \cdot \beta_j = s_{\gamma_j} \cdot \gamma_j = -\gamma_j \in \Phi^{-},$$

therefore Lemma 4.2 gives us a standard expression of $w_{i,j-1}$, hence of $w_{i,n}$, beginning with s_{γ_i} . The same situation occurs if $\beta_i \in \Phi^{-}$. However, this prevents the given expression of w from being standard, contradicting the hypothesis. Thus all β_i are positive and distinct. Moreover, Lemma 2.2 implies that $\beta_i \in \Phi[s_{\gamma_i} w_{i,n}] \subseteq \Phi[w]$. This completes the first claim of this proof.

From now, we prove the other inequality $|\Phi^{\perp[x]}[w]| \leq n$ by induction on n . First, Lemma 4.1 says that any non-loop generator of C , hence the u_n , lies in the groupoid Y . This proves the claim for $n = 0$ and allows us to assume that $u_n = 1$ without loss of generality. Now since $\Phi^{\perp[x]}[s_{\gamma_n}] = \{\gamma_n\}$ (see Lemma 4.1), s_{γ_n} maps $\Phi^{\perp[x]}[w] \setminus \{\gamma_n\}$ into the set $\Phi^{\perp[x]}[w_{0,n-1}]$ with cardinality not larger than $n - 1$ (the induction assumption), proving the desired inequality. Hence the proof of Proposition 4.3 is concluded. $\square$

The next result shows a further remarkable property of C , enhancing Theorem 3.5(3). Note that the corresponding property of the fundamental relations of Coxeter groups plays a crucial role in the solution of the word problem in Coxeter groups; see e.g., [2, Theorem 3.3.1]. The proof is analogous to the case of Coxeter groups again.

Proposition 4.4. *Let $w \in C$ be any element.*

1. *Any two standard expressions of w can be converted to each other by using the generalized braid moves only.*
2. *Any expression of w can be converted to a given standard expression of w by using the generalized braid moves and cancellations of subwords of the form $w_{\xi} w_{\varphi(\xi)}$ only, not using insertions of subwords $w_{\xi} w_{\varphi(\xi)}$.*

Proof. For the claim 1, we proceed the proof by induction on $\ell(w)$, the case $\ell(w) = 0$ being trivial. Suppose that $\ell(w) > 0$, and let $w_{x_1}^{s_1} \cdots w_{x_n}^{s_n}$ and $w_{y_1}^{t_1} \cdots w_{y_m}^{t_m}$ be two standard expressions of w , hence $x_n = y_m$. Then we have $w \cdot \alpha_{s_n} \in \Phi^{-}$ by the fact $w_{x_n}^{s_n} \cdot \alpha_{s_n} \in \Phi^{-}$ and Lemma 2.2, and $w \cdot \alpha_{t_m} \in \Phi^{-}$ similarly. Now the combination of Lemma 3.2 and Theorem 3.5(1)(2) implies that $J = [x_n]_{\sim \{s_n, t_m\}}$ is of finite type and there are two standard expressions of w of the forms $u c_1 \cdots c_r$ and $u c'_1 \cdots c'_r$, where $c_1 \cdots c_r$ and $c'_1 \cdots c'_r$ are the standard expressions of $w_0(J) w_0(J \setminus \{s_n, t_m\}) \in C$ ending with $c_r = w_{x_n}^{s_n}$ and $c'_r = w_{y_m}^{t_m}$, respectively. Then we have

$$w_{x_1}^{s_1} \cdots w_{x_n}^{s_n} \xrightarrow{GBMs} u c_1 \cdots c_r \xrightarrow{GBM} u c'_1 \cdots c'_r \xrightarrow{GBMs} w_{y_1}^{t_1} \cdots w_{y_m}^{t_m}$$

as desired, where the first and the third transformations come from the induction assumption and the second one is the definition of the GBM.

For the claim 2, let $w_{\xi_1} \cdots w_{\xi_n}$ be an expression of w . We show the claim by induction on the total length of the w_{ξ_i} , the case of length zero being trivial. Owing to the claim 1, we may assume that this expression is not standard. Take the last index i such that $w_{\xi_1} \cdots w_{\xi_{i-1}}$ is standard. Then by Lemma 4.2, $w_{\xi_1} \cdots w_{\xi_{i-1}}$ admits another standard expression of the form $w_{\zeta_1} \cdots w_{\zeta_m} w_{\varphi(\xi_i)}$, which can be reached from $w_{\xi_1} \cdots w_{\xi_{i-1}}$ by GBMs only (apply the claim 1). Thus we have

$$w_{\xi_1} \cdots w_{\xi_n} \xrightarrow{GBMs} w_{\zeta_1} \cdots w_{\zeta_m} w_{\varphi(\xi_i)} w_{\xi_i} \cdots w_{\xi_n} \xrightarrow{cancel} w_{\zeta_1} \cdots w_{\zeta_m} w_{\xi_{i+1}} \cdots w_{\xi_n}$$

and the total length decreases through the transformation. Hence the induction works and the claim follows. $\square$

Remark 4.5. As shown in Theorem 3.5(2), both terms of a generalized braid move have the same number of generators, the same number of loop generators (see Proposition 4.3) and the same total length of generators contained. Thus none of those quantities increases in a transformation appearing in Proposition 4.4. As a result, any expression of $w \in C$ contains at least $\text{lp}(w)$ loop generators.

4.2 The factorization of C

The previous results enable us to deduce the following properties. Recall from Section 3.1 that $W^{\perp[x]}$ is normal in $C_{x,x}$ and $W^{\perp[x]} \cap Y_{x,x} = 1$.

Theorem 4.6. 1. *We have*

$$C_{y,x} = W^{\perp[y]} \cdot Y_{y,x} = Y_{y,x} \cdot W^{\perp[x]} \text{ for all } x, y \in \mathcal{V}(C) .$$

Hence $C_{x,x}$ admits a semidirect product decomposition $C_{x,x} = W^{\perp[x]} \rtimes Y_{x,x}$.

2. *The groupoid Y is generated by all the non-loop generators of C . We have*

$$Y_{y,x} = \{w \in C_{y,x} \mid \text{lp}(w) = 0\} \text{ for all } x, y \in \mathcal{V}(C) .$$

Moreover, Proposition 4.4 holds also for Y under the modification that the generalized braid moves are restricted to the ones containing no loop generators.

3. *The length of an element $w \in W^{\perp[x]}$ with respect to the generating set $R^x = S(\Phi^{\perp[x]})$ of $W^{\perp[x]}$ is equal to $\text{lp}(w)$, and we have*

$$R^{[x]} = \{ws_{\gamma}w^{-1} \mid s_{\gamma} \in C_{y,y} \text{ and } w \in Y_{x,y} \text{ for some } y \in \mathcal{V}(C)\} . \quad (4.4)$$

Moreover, $w \in W^{\perp[x]}$ admits an expression $(w_1 s_{\gamma_1} w_1^{-1}) \cdots (w_n s_{\gamma_n} w_n^{-1})$ with $n = \text{lp}(w)$ such that the expression obtained from it by replacing every $w_1, w_i^{-1} w_{i+1}, w_n^{-1}$ with its standard expression is also standard.

4. *If $w \in Y_{y,x}$, then the map $u \mapsto wuw^{-1}$ is an isomorphism of Coxeter systems from $(W^{\perp[x]}, R^{[x]})$ to $(W^{\perp[y]}, R^{[y]})$.*

Proof. First we note the following fact used in the proof. If $w \in C$ admits an expression of the form $u_0 s_{\gamma_1} u_1 s_{\gamma_2} \cdots u_{n-1} s_{\gamma_n} u_n$, where the s_{γ_i} are loop generators and the u_i contain no loop generators, then

$$w = (w_1 s_{\gamma_1} w_1^{-1} \cdots w_n s_{\gamma_n} w_n^{-1}) w_{n+1} = (s_{w_1 \cdot \gamma_1} \cdots s_{w_n \cdot \gamma_n}) w_{n+1} \quad (4.5)$$

where we put $w_i = u_0 u_1 \cdots u_{i-1}$ for each $1 \leq i \leq n+1$.

For the claim 1, it suffices to show the first part of the claim, in particular the first equality, since then the second equality follows by taking the inverse. This is done by using the transformation (4.5) and Lemma 4.1 (indeed, if $w \in C_{y,x}$, then $w_i \cdot \gamma_i \in \Phi^{\perp[y]}$ and $w_{n+1} \in Y_{y,x}$).

For the claim 2, the first and the third parts follow from the second part, Theorem 3.5(1) and Remark 4.5. Moreover, the second one is deduced from Proposition 4.3.

For the claim 3, note that the claim on the length of w follows from Theorem 2.3(5) and Proposition 4.3. Moreover, any element u of the right-hand side of (4.4) satisfies that $\text{lp}(u) \leq 1$ (see Remark 4.5) and $1 \neq u \in W^{\perp[x]}$, hence $\text{lp}(u) = 1$ and $u \in R^{[x]}$ by the above claim. Now take a standard expression $u_0 s_{\gamma_1} u_1 s_{\gamma_2} \cdots u_{n-1} s_{\gamma_n} u_n$ of $w \in W^{\perp[x]}$ in C of the above form, hence $n = \text{lp}(w)$. Then in the equality (4.5), we have $w_{n+1} = 1$ since $W^{\perp[x]} \cap Y_{x,x} = 1$, while $w_1 = u_0$ and $w_i^{-1} w_{i+1} = u_i$ for all $1 \leq i \leq n$. Hence (4.5) is the desired expression of w . This argument also implies that the set $R^{[x]} = \{w \in W^{\perp[x]} \mid \text{lp}(w) = 1\}$ is contained in the right-hand side of (4.4). Hence the claim holds.

Finally, the claim 4 is a corollary of the claim 3. Hence Theorem 4.6 holds. $\square$

Remark 4.7. It is shown in [5] that the vertex group $G_I = G_{I,I}$ of the groupoid G admits a similar decomposition $G_I = \tilde{N}_I \rtimes M_I$ into a Coxeter group $\tilde{N}_I$ and a vertex group M_I of a groupoid M , where M and $\tilde{N}_I$ are (by definition) generated by non-loop generators of G , and by conjugates of loop generators of G by elements of M , respectively. However, in contrast with the case of C , the properties of $W^{\perp I}$ and Y are not inherited immediately from $\tilde{N}_I$ and M , since a lift of a loop generator of G may be a non-loop generator of C and not all the generators of $W^{\perp I}$ are lifts of those of $\tilde{N}_I$. More precisely, it can be shown that $W^{\perp I}$ is a normal reflection subgroup of the Coxeter group $\tilde{N}_I$, but *not* a parabolic one; hence the structure of $W^{\perp I}$ is still hard to describe even if that of $\tilde{N}_I$ is clear.

4.3 The factor Y

In this and the following subsections, we study the structure of the two factors $W^{\perp I}$ and Y_I more precisely. First, we have the following result:

Proposition 4.8. *Each vertex group $Y_{x,x}$ of Y is torsion-free.*

Proof. If $w \in Y_{x,x}$ has finite order, then Proposition 2.8 gives us a decomposition $w = s_{\beta_1} \cdots s_{\beta_r}$ such that each s_{β_j} fixes $\Pi_{[x]}$ pointwise, therefore we have $\beta_j \in \Phi^{\perp[x]}$ and $w \in Y_{x,x} \cap W^{\perp[x]} = 1$, implying $w = 1$ as desired. $\square$

This yields the following analogy of Lemma 3.6 (see Section 3.4 for the definitions):

Corollary 4.9. *Any circular tour has order 1. Moreover, for a path p in the graph $\mathcal{C}$, the followings are equivalent:*

1. p is a circular tour;
2. $p = (c'_1 \cdots c'_r)^{-1} c_1 \cdots c_r$ as paths in $\mathcal{C}$ for a generalized braid move $c_1 \cdots c_r \rightsquigarrow c'_1 \cdots c'_r$ with loop number 0.

Proof. By definition, any circular tour p represents an element of finite order in some $Y_{x,x}$ that is a torsion-free group (see Proposition 4.8), hence the element must be identity. The second claim is deduced by the same argument as Lemma 3.6. $\square$

The combination of this corollary and Theorem 4.6(2) yields the next description of the groupoid Y . Define a complex $\mathcal{Y}$ (see Section 2.1 for terminology) such that its 1-skeleton $\mathcal{Y}^1$ is the subgraph of $\mathcal{C}$ obtained by deleting all the loops, and $\mathcal{Y}$ has a 2-cell with boundary $\mathcal{G}$ for each cycle $\mathcal{G}$ as in the definition of circular tours (see Section 3.4). Note that every path in $\mathcal{Y}$ represents an element of C .

Theorem 4.10. *The groupoid Y is naturally isomorphic to the fundamental groupoid $\pi_1(\mathcal{Y}; *, *)$ of the complex $\mathcal{Y}$. Hence each $Y_{x,x}$ is also isomorphic to the fundamental group $\pi_1(\mathcal{Y}; x)$ of $\mathcal{Y}$ at x .*

Note that the groupoid C also admits a similar description in terms of a certain complex with $\mathcal{C}$ being the 1-skeleton, though $\mathcal{C}$ involves a loop in general and the boundary of a 2-cell is not always a simple closed path any longer.

For a further description, we fix a maximal tree $\mathcal{T}$ in the connected graph $\mathcal{Y}^1$. For $y, z \in \mathcal{V}(C)$, let $p_{z,y}$ denote the unique non-backtracking path in $\mathcal{T}$ from y to z . Note that $p_{z,y}p_{y,x} = p_{z,x}$ and $p_{y,z} = p_{z,y}^{-1}$ hold in $\pi_1(\mathcal{Y}^1; *, *)$. Moreover, put

$$q_{(x)} = p_{x,z}qp_{y,x} \quad \text{for } x \in \mathcal{V}(C) \text{ and a path } q \text{ in } \mathcal{Y}^1 \text{ from } y \text{ to } z, \quad (4.6)$$

which is the extension of the path q to a vertex x along the tree $\mathcal{T}$. Note that $(e_{\xi_n} \cdots e_{\xi_1})_{(x)} = (e_{\xi_n})_{(x)} \cdots (e_{\xi_1})_{(x)}$ in $\pi_1(\mathcal{Y}^1; x)$. Then a theorem in combinatorial group theory (see e.g., [7, Theorem 5.17]) yields the following presentation of $\pi_1(\mathcal{Y}; x)$:

Theorem 4.11. *The group $\pi_1(\mathcal{Y}; x)$ admits a presentation with generators given by*

$$(w_y^s)_{(x)} = p_{x,z}w_y^s p_{y,x}, \quad \text{with } w_y^s \in E(\mathcal{Y}) \text{ and } z = w_y^s \cdot y,$$

where $E(\mathcal{Y})$ denotes the set of the oriented edges of $\mathcal{Y}$, and fundamental relations given by

- $(w_\xi)_{(x)}(w_{\varphi(\xi)})_{(x)} = 1$ for every $w_\xi \in E(\mathcal{Y})$;
- $(w_\xi)_{(x)} = 1$ for every $w_\xi \in E(\mathcal{Y})$ contained in the tree $\mathcal{T}$;
- $(e_{\xi_n})_{(x)} \cdots (e_{\xi_1})_{(x)} = 1$ for a boundary $e_{\xi_n} \cdots e_{\xi_1}$ of each 2-cell of $\mathcal{Y}$.

Example 4.12. Here we consider the case $|I| = 1$ examined by Brink [4]; hence $S^{(\Lambda)} = S$ and $I = \{x_I\}$. Then for $x \in S$ and $s \in S \setminus \{x\}$, the set $\{x\}_{\sim_s}$ is of finite type and $w_x^s \cdot x \neq x$ if and only if $m_{x,s}$ is odd. Thus the 1-skeleton $\mathcal{Y}^1$ is the connected component of the *odd Coxeter graph* Γ^{odd} of (W, S) containing x_I used in Brink's description of $Z_W(x_I)$, where Γ^{odd} is (by definition) the subgraph of Γ obtained by removing the edges with non-odd labels. Moreover, the acyclicity of Coxeter graphs of finite type implies that this $\mathcal{Y}$ has no 2-cell and $\mathcal{Y} = \mathcal{Y}^1$. Hence our result agrees with the result of [4] for this special case.

4.4 The factor $W^{\perp I}$

First, for $x \in \mathcal{V}(C)$, define

$$\mathcal{R}_x = \{(w, u) \mid w \in Y_{x,x} \text{ and } u \text{ is a loop generator of } C\}$$

and

$$r_x(w, u) = wu_{(x)}w^{-1} \text{ for } (w, u) \in \mathcal{R}_x \text{ (see (4.6) for the notation),}$$

hence the generating set $R^{[x]}$ of $W^{\perp[x]}$ consists of all the $r_x(w, u)$ with $(w, u) \in \mathcal{R}_x$ (see Theorem 4.6(3)). The subscripts ' x ' are omitted when $x = x_I$. The pair (w, u) is also denoted by (w, γ)

when $u = s_\gamma$. To study relations between the generators $r_x(w, u)$, for integer $1 \leq k < \infty$, we define a relation $\overset{k}{\sim}$ on $\mathcal{R}_x$ by

$$(w, \gamma) \overset{k}{\sim} (wq_{(x)}, \beta) \text{ if there is a shuttling tour of order } k \text{ of the form } s_\gamma q s_\beta q^{-1}$$

(see Section 3.4 for the terminology). Note that the relation $\overset{k}{\sim}$ is symmetric, and the product $r_x(w, \gamma)r_x(u, \beta)$ of two generators of $W^{\perp[x]}$ has order k if $(w, \gamma) \overset{k}{\sim} (u, \beta)$. Let $\sim$ denote the transitive closure of $\overset{1}{\sim}$, which is an equivalence relation.

The next result describes the structure of the group $W^{\perp[x]}$ in terms of those relations. Note that the first claim of this result can be deduced directly from Theorem 3.5(3) in a similar way, without Proposition 4.4 that is essential in the proof of the second claim.

Theorem 4.13. *The group $W^{\perp[x]}$ admits a presentation with generators given by*

$$r_x(w, \gamma), \text{ with } (w, \gamma) \in \mathcal{R}_x$$

and fundamental relations given by

- $r_x(w, \gamma)^2 = 1$ for every $(w, \gamma) \in \mathcal{R}_x$;
- $(r_x(w, \gamma)r_x(u, \beta))^k = 1$ for every pair $(w, \gamma) \overset{k}{\sim} (u, \beta)$ with $1 \leq k < \infty$.

Moreover, any expression of an element of $W^{\perp[x]}$ (with respect to these generators) can be converted into an arbitrarily given shortest expression of the element (with respect to these generators) by using the following two kinds of transformations:

- $r_x(w, \gamma)^2 \rightsquigarrow 1$, corresponding to the first relation above;
- $r_x(w, \gamma)r_x(u, \beta)r_x(w, \gamma) \cdots \rightsquigarrow r_x(u, \beta)r_x(w, \gamma)r_x(u, \beta) \cdots$ (where both terms consist of k generators), corresponding to the second relation above.

Proof. Owing to Theorem 4.6(3), it suffices to prove that any expression of an element of $W^{\perp[x]}$ can be converted into a given shortest expression of the form given in Theorem 4.6(3) in the above way, since the above transformations of second type are invertible. In this proof, we write $w \equiv u$ if two expressions w and u represent the same element, distinguishing from the case $w = u$ where the expressions w and u themselves are equal.

To prove the above claim, write the given original expression as

$$w_1 l_1 w_1^{-1} w_2 l_2 w_2^{-1} \cdots w_n l_n w_n^{-1}, \text{ where } (w_i, l_i) \in \mathcal{R}_x. \quad (4.7)$$

Put $w_0 = 1$ and $w_{n+1} = 1$. Then, owing to the assumption on the property of the target shortest expression, the combination of Proposition 4.4 and Theorem 4.6(2) implies the following fact: To convert (4.7) into the target expression, where both of the original and the target expressions are regarded as those in C , the following three kinds of transformations are enough and an insertion of any subword $s_\beta s_\beta$ is *not* required:

- (T1) a cancellation of a subword $s_\beta s_\beta$, with s_β a loop generator;
- (T2) a GBM that contains a loop generator;
- (T3) a transformation $u \rightsquigarrow u'$, where u and u' contain no loop generators and $u \equiv u'$ in Y .

Now we proceed the proof by induction on the total number N of transformations T1 and T2 in the conversion process. Note that, in the case $N = 0$, the original and the target expressions with respect to the generators of $W^{\perp[x]}$ are already equal, hence we have nothing to do.

In the conversion process of the general case, we apply (possibly no) T3 transformations to the expression (4.7) before applying the first T1 or T2 transformation, obtaining an expression

$$u_1 l_1 u_2 l_2 u_3 \cdots u_n l_n u_{n+1}, \text{ where the } u_i \text{ contain no loop generators and } u_i \equiv w_{i-1}^{-1} w_i. \quad (4.8)$$

First we consider the case that T1 is then applied to (4.8). If the generators l_{i-1} and l_i with $2 \leq i \leq n$ are cancelled by the T1, then u_i should be empty (hence $w_{i-1} \equiv w_i$) and $l_{i-1} = l_i$. Now the T1 results in the expression

$$u_1 l_1 u_2 \cdots u_{i-2} l_{i-2} u_{i-1} u_{i+1} l_{i+1} u_{i+2} \cdots u_n l_n u_{n+1}. \quad (4.9)$$

Moreover, since now $u_{i-1} u_{i+1} \equiv w_{i-2}^{-1} w_{i+1}$, (4.9) is convertible by a sequence of T3 transformations into

$$w_1 l_1 w_1^{-1} \cdots w_{i-2} l_{i-2} w_{i-2}^{-1} w_{i+1} l_{i+1} w_{i+1}^{-1} \cdots w_n l_n w_n^{-1}, \quad (4.10)$$

which is also obtained from (4.7) by the transformation $r_x(w_i, l_i)^2 \rightsquigarrow 1$ in the statement (recall that now $w_{i-1} \equiv w_i$ in Y and $l_{i-1} = l_i$, hence $r_x(w_{i-1}, l_{i-1}) = r_x(w_i, l_i)$). Now the total number of T1 and T2 transformations required in a conversion of (4.10) into the target expression is the same as that for (4.9), which is $N - 1$, hence the claim follows by the induction assumption.

From now, we consider the other case that T2 is applied to (4.8) instead of T1. Then the subword of (4.8) on which the GBM acts can be written as $bl_i u_{i+1} \cdots u_j l_j c$ for some $1 \leq i < j \leq n$, where we decomposed u_i and u_{j+1} as $u_i = ab$ and $u_{j+1} = cd$. Here we only consider the case that the loop number $k = j - i + 1$ of this GBM is odd, since the other case is similar. Then Lemma 3.6 implies that, by the shape of shuttling tours (see Figure 1), there is a shuttling tour $p = lqLq^{-1}$ of order k , where l and L are loop generators, such that

$$l_i = l_{i+2} = \cdots = l_j = l, \quad l_{i+1} = l_{i+3} = \cdots = l_{j-1} = L, \\ u_{i+1} = u_{i+3} = \cdots = u_{j-1} = q, \quad u_{i+2} = u_{i+4} = \cdots = u_j = q^{-1}$$

and the expression q admits two decompositions of the form $q = cc'$ and $q = b^{-1}b'^{-1}$. By these and the relations $u_m \equiv w_{m-1}^{-1} w_m$, we have $w_i \equiv w_{i+2} \equiv \cdots \equiv w_j$ and $w_{i+1} \equiv w_{i+3} \equiv \cdots \equiv w_{j-1}$ in Y . Now this GBM, whose loop number is k , is of the form

$$bl_i u_{i+1} \cdots u_j l_j c = b(lqLq^{-1})^{(k-1)/2} l c \overset{GBM}{\rightsquigarrow} b'^{-1} L q^{-1} (lqLq^{-1})^{(k-3)/2} l q L c'^{-1},$$

which converts (4.8) into

$$u_1 l_1 \cdots u_{i-1} l_{i-1} a \left(b'^{-1} L q^{-1} (lqLq^{-1})^{(k-3)/2} l q L c'^{-1} \right) d l_{j+1} u_{j+2} \cdots l_n u_{n+1}. \quad (4.11)$$

On the other hand, the shuttling tour p defines a relation $(w_i, l) \overset{k}{\sim} (w_i q, L)$, while $w_i q = w_i u_{i+1} \equiv w_{i+1}$. Thus we have

$$(4.7) \equiv w_1 l_1 w_1^{-1} \cdots w_{i-1} l_{i-1} w_{i-1}^{-1} (w_i l w_i^{-1} w_{i+1} L w_{i+1}^{-1})^{(k-1)/2} w_i l w_i^{-1} \\ \cdot w_{j+1} l_{j+1} w_{j+1}^{-1} \cdots w_n l_n w_n^{-1} \\ \rightsquigarrow w_1 l_1 w_1^{-1} \cdots w_{i-1} l_{i-1} w_{i-1}^{-1} (w_{i+1} L w_{i+1}^{-1} w_i l w_i^{-1})^{(k-1)/2} w_{i+1} L w_{i+1}^{-1} \\ \cdot w_{j+1} l_{j+1} w_{j+1}^{-1} \cdots w_n l_n w_n^{-1}, \quad (4.12)$$

where the left-hand and the middle terms coincide with each other as expressions with respect to the generators of $W^{\perp[x]}$ given in the statement, and the last transformation comes from a transformation of second type given in the statement concerning the generators $r_x(w_i, l)$ and $r_x(w_{i+1}, L)$. Moreover, we have

$$w_{i-1}^{-1}w_{i+1} \equiv u_i u_{i+1} \equiv ab'^{-1}, \quad w_{i+1}^{-1}w_i \equiv q^{-1}, \quad w_{i+1}^{-1}w_{j+1} \equiv u_j u_{j+1} \equiv c'^{-1}d, \quad ,$$

therefore the right-hand side of (4.12) is convertible into (4.11) by a sequence of T3 transformations. Thus the total number of T1 and T2 transformations required in a conversion of the right-hand side of (4.12) into the target expression is the same as that for (4.11), which is $N - 1$, therefore the claim follows from the induction assumption. Hence the proof of Theorem 4.13 is concluded. $\square$

This theorem yields the main result of this subsection, as follows:

Theorem 4.14. *We have $r_x(w, \gamma) = r_x(u, \beta)$ if and only if $(w, \gamma) \sim (u, \beta)$. Moreover, $r_x(w, \gamma)r_x(u, \beta)$ has order k with $2 \leq k < \infty$ if and only if there exist (w', γ') and (u', β') in $\mathcal{R}_x$ such that $(w, \gamma) \sim (w', \gamma') \stackrel{k}{\sim} (u', \beta') \sim (u, \beta)$.*

Proof. Note that both of the two “if” parts follow from the definition. From now, we prove the two “only if” parts. First, note that if $(w, \gamma) \sim (w', \gamma')$, $(u, \beta) \sim (u', \beta')$, $(w, \gamma) \stackrel{k}{\sim} (u, \beta)$ and $(w', \gamma') \stackrel{\ell}{\sim} (u', \beta')$, then we have $k = \ell$, since now k and ℓ are the order of the same element $r_x(w, \gamma)r_x(u, \beta) = r_x(w', \gamma')r_x(u', \beta')$. Now Theorem 4.13 yields a presentation of $W^{\perp[x]}$ with generators given by

$$r_x(\overline{w, \gamma}) = r_x(w, \gamma), \quad \text{where } \overline{w, \gamma} \text{ denotes an equivalence class of } \sim$$

and fundamental relations given by

- $r_x(\overline{w, \gamma})^2 = 1$ for every generator $r_x(\overline{w, \gamma})$,
- $(r_x(\overline{w, \gamma})r_x(\overline{u, \beta}))^k = 1$ for every pair $(w, \gamma) \stackrel{k}{\sim} (u, \beta)$.

The above remark shows that these data form a Coxeter presentation of the Coxeter group $W^{\perp[x]}$. Thus for two (w, γ) and (u, β) in $\mathcal{R}_x$, the product $r_x(w, \gamma)r_x(u, \beta)$ has order $1 \leq k < \infty$ if and only if the relation “ $(r_x(\overline{w, \gamma})r_x(\overline{u, \beta}))^k = 1$ ” appears as one of the above fundamental relations. Hence the claim follows. $\square$

To study the relation $\sim$ further, we introduce an auxiliary graph $\mathcal{I}$ as follows. The vertices of $\mathcal{I}$ are the loop generators s_γ of C , or equivalently the loops in the graph $\mathcal{C}$. The edges of $\mathcal{I}$ from a vertex s_β to a vertex s_γ are the paths q in $\mathcal{Y}^1$ such that $s_\gamma q s_\beta q^{-1}$ is a shuttling tour of order one (note that such a path q must be nonempty). Let $\iota : \pi_1(\mathcal{I}; *, *) \rightarrow \pi_1(\mathcal{Y}^1; *, *)$ be the groupoid homomorphism that maps a vertex s_γ of $\mathcal{I}$ to $x \in \mathcal{V}(C)$ with $s_\gamma \in C_{x,x}$, and maps an edge q of $\mathcal{I}$ to a path q in $\mathcal{Y}^1$.

Remark 4.15. Note that the induced homomorphism $\iota : \pi_1(\mathcal{I}; s_\gamma) \rightarrow \pi_1(\mathcal{Y}^1; x)$ between fundamental groups, where $s_\gamma \in C_{x,x}$, is *injective*. Indeed, if a cancellation of two consecutive edges ee^{-1} occurs in a path $\iota(q_n \cdots q_2 q_1)$, then it should occur between the end of some $\iota(q_i)$ and the beginning of $\iota(q_{i+1})$, forcing $q_{i+1} = q_i^{-1}$ by definition of $\mathcal{I}$ and reducing $q_n \cdots q_1$ to a shorter path $q_n \cdots q_{i+2} q_{i-1} \cdots q_1$ in $\mathcal{I}$. This argument implies the injectivity of ι .

Now Theorem 4.14 implies that $r_x(w, \gamma) = r_x(u, \beta)$ if and only if there is a path $\tilde{q}$ in $\mathcal{I}$ from s_β to s_γ such that $u = w\iota(\tilde{q})_{(x)}$. Hence $r_x(w, \gamma) = r_x(u, \gamma)$ if and only if $(w^{-1}u)_{(y)} \in \iota(\pi_1(\mathcal{I}; s_\gamma))$ where $s_\gamma \in C_{y,y}$.

By this formulation, we obtain the following observation about the *finite part* of the Coxeter group $W^{\perp I}$, denoted by $W^{\perp I}_{\text{fin}}$, which we define as the product of all the irreducible components of $W^{\perp I}$ of finite type. We prepare some temporary notations. For a fixed loop generator $s_\gamma \in C_{x,x}$, put $P = \pi_1(\mathcal{I}; s_\gamma)$ and let N denote the kernel of the natural projection $\pi_1(\mathcal{Y}^1; x) \twoheadrightarrow \pi_1(\mathcal{Y}; x)$ where the image of c is denoted by $\bar{c}$. Let $\bar{\iota}$ denote the composition $\pi_1(\mathcal{I}; *, *) \xrightarrow{\iota} \pi_1(\mathcal{Y}^1; *, *) \twoheadrightarrow \pi_1(\mathcal{Y}; *, *)$ of the above map ι followed by the natural projection. Let Ab denote the abelianization map $\pi_1(\mathcal{Y}^1; x) \twoheadrightarrow \text{Ab}(\pi_1(\mathcal{Y}^1; x))$.

Theorem 4.16. *Under the above notations, suppose that $|S| < \infty$ and the subgroup $\text{Ab}(\iota(P)N)$ has infinite index in $\text{Ab}(\pi_1(\mathcal{Y}^1; x))$. Then for any $w \in Y_I$, the vertex $r(w, \gamma)$ of the Coxeter graph of $W^{\perp I}$ is adjacent to an edge with label ∞ , hence $r(w, \gamma) \notin W^{\perp I}_{\text{fin}}$.*

In particular, the second assumption is satisfied if $\text{rk}(P) + n < \text{rk}(\pi_1(\mathcal{Y}^1; x))$, where $\text{rk}(F)$ denotes the rank of a free group F and n is the number of 2-cells of $\mathcal{Y}$.

Proof. To prove the first claim, we suppose that $r(w, \gamma)$ is not adjacent to an edge with label ∞ in the Coxeter graph of $W^{\perp I}$, and give a decomposition of $\text{Ab}(\pi_1(\mathcal{Y}^1; x))$ into a finite number of cosets modulo $\text{Ab}(\iota(P)N)$, yielding a contradiction. Theorem 4.6(4) allows us to assume without loss of generality that $w = 1$. Then for any $a \in \pi_1(\mathcal{Y}^1; x)$, we have either $r(1, \gamma) = r(\bar{a}_{(x_I)}, \gamma)$, or $r(1, \gamma)r(\bar{a}_{(x_I)}, \gamma)$ has order $2 \leq k < \infty$, by the above hypothesis of the proof.

In the second case, Theorem 4.14 implies the existence of an element $b \in \pi_1(\mathcal{Y}^1; x_I)$ and a shuttling tour $s_\beta q s_{\beta'} q^{-1}$ of order k such that

$$(1, \gamma) \sim (\bar{b}, \beta) \stackrel{k}{\sim} (\bar{b}(\bar{q})_{(x_I)}, \beta') \sim (\bar{a}_{(x_I)}, \gamma) . \quad (4.13)$$

Now the hypothesis $|S| < \infty$ implies that only a finite number of shuttling tours, say $s_{\beta_i} q_i s_{\beta'_i} q_i^{-1}$ with $1 \leq i \leq r < \infty$, appear in this manner. Let the above-mentioned shuttling tour be the j -th one. Then (4.13) implies that $\bar{b} = \bar{\iota}(\tilde{q})_{(x_I)}$ for some $\tilde{q} \in \pi_1(\mathcal{I}; s_\gamma, s_{\beta_j})$, and $\bar{a}_{(x_I)} = \bar{b}(\bar{q}_j)_{(x_I)} \bar{\iota}(\tilde{q}')_{(x_I)}$ for some $\tilde{q}' \in \pi_1(\mathcal{I}; s_{\beta'_j}, s_\gamma)$. Thus we have $\bar{a}_{(x_I)} = \overline{(\iota(\tilde{q})q_j \iota(\tilde{q}'))_{(x_I)}}$. Since $s_\gamma \in C_{x,x}$ and $a \in \pi_1(\mathcal{Y}^1; x)$, it follows that

$$\bar{a} = \overline{\iota(\tilde{q})q_j \iota(\tilde{q}')} \in \overline{\iota(P)\iota(\tilde{p}_j)q_j \iota(\tilde{p}'_j)\iota(P)}$$

for some $\tilde{p}_j \in \pi_1(\mathcal{I}; s_\gamma, s_{\beta_j})$ and $\tilde{p}'_j \in \pi_1(\mathcal{I}; s_{\beta'_j}, s_\gamma)$ (fixed for each j), therefore $a \in \iota(P)c_j \iota(P)N$ where we put $c_j = \iota(\tilde{p}_j)q_j \iota(\tilde{p}'_j) \in \pi_1(\mathcal{Y}^1; x)$.

On the other hand, in the first case $r(1, \gamma) = r(\bar{a}_{(x_I)}, \gamma)$, we have $\bar{a} \in \bar{\iota}(P)$ and $a \in \iota(P)c_0 \iota(P)N$, where we put $c_0 = 1 \in \pi_1(\mathcal{Y}^1; x)$.

Summarizing, we have $a \in \iota(P)c_i \iota(P)N$ for some $0 \leq i \leq r$. Since a was arbitrarily chosen, this means that $\pi_1(\mathcal{Y}^1; x) = \bigcup_{i=0}^r \iota(P)c_i \iota(P)N$, therefore we have

$$\text{Ab}(\pi_1(\mathcal{Y}^1; x)) = \bigcup_{i=0}^r \text{Ab}(\iota(P))\text{Ab}(c_i)\text{Ab}(\iota(P))\text{Ab}(N) = \bigcup_{i=0}^r \text{Ab}(c_i)\text{Ab}(\iota(P)N) .$$

This is the desired decomposition of $\text{Ab}(\pi_1(\mathcal{Y}^1; x))$ into finitely many cosets modulo $\text{Ab}(\iota(P)N)$. Hence the first claim holds.

For the second claim, let $\tilde{g}_1, \dots, \tilde{g}_r$ (where $r = \text{rk}(P)$) be free generators of P and $c_1, \dots, c_n$ the boundaries of the 2-cells of $\mathcal{Y}$. Then N is generated by the conjugates of the $(c_i)_{(x)}$,

therefore $\text{Ab}(\iota(P)N)$ is generated by the $\text{Ab}(\iota(\tilde{g}_i))$ and $\text{Ab}((c_j)_{(x)})$. By the hypothesis, the total number $r + n$ of these generators is less than $\text{rk}(\pi_1(\mathcal{Y}^1; x))$ that is equal to the rank of the free abelian group $\text{Ab}(\pi_1(\mathcal{Y}^1; x))$, implying that the subgroup $\text{Ab}(\iota(P)N) \subseteq \text{Ab}(\pi_1(\mathcal{Y}^1; x))$ has infinite index, as desired. Hence Theorem 4.16 holds. $\square$

On the other hand, for a general S , we have the following result. For any $J \subseteq S$, put

$$J^\perp = \{s \in S \setminus J \mid s \text{ is not adjacent to } J \text{ in the graph } \Gamma\} . \quad (4.14)$$

Proposition 4.17. *Let $s_\gamma = w_x^s$ be a loop generator of C , $w \in Y_I$ and $J \subseteq S$ a finite subset such that $[x]_{\sim_s} \subseteq J$ and $[x] \setminus J \subseteq J^\perp$, hence $s_\gamma \in R^{J, [x] \cap J}$. Then we have $r(w, \gamma) \notin W_{\text{fin}}^{\perp [x]}$ whenever $s_\gamma \notin W_J^{\perp [x] \cap J}$.*

Proof. First, Theorem 4.6(4) allows us to assume that $w = 1$ and $x = x_I$, therefore $r(w, \gamma) = s_\gamma$. Now since $[x] \setminus J \subseteq J^\perp$, we have $\Phi_J^{\perp [x]} = \Phi_J^{\perp [x] \cap J}$, hence $\Pi^{J, [x] \cap J} = \Pi^{J, [x]} \subseteq \Pi^{[x]}$. This implies that the Coxeter system $(W_J^{\perp [x] \cap J}, R^{J, [x] \cap J})$ is a subsystem of $(W^{\perp [x]}, R^{[x]})$. Hence if $s_\gamma \notin W_J^{\perp [x] \cap J}$, then the irreducible component of $W^{\perp [x]}$ containing $s_\gamma = r(w, \gamma)$ contains the infinite irreducible component of $W_J^{\perp [x] \cap J}$ containing s_γ , implying that $r(w, \gamma) \notin W_{\text{fin}}^{\perp [x]}$ as desired. $\square$

At the last of this subsection, we explain how to compute the order of a shuttling tour in an individual case. We fix a shuttling tour $s_\gamma u s_\beta u^{-1}$ in a connected component $\mathcal{G}$ of $\mathcal{C}(J)$ (recall the definition in Section 3.4) with $\beta, \gamma \in \Phi^+$, and put $s_\beta = w_x^s$, $s_\gamma = w_y^t$ and $J = [x] \cup \{s, s'\} = [y] \cup \{t, t'\}$. Put $K = J_{\sim(J \setminus [y])}$, which is of finite type. Now by applying our arguments above to $W_J^{\perp [y]}$ instead of $W^{\perp [y]}$, it follows that $R^{J, [y]} = \{s_\gamma, u s_\beta u^{-1} = s_{u \cdot \beta}\}$ and, since $u \cdot \beta \in \Phi^+$, we have $\Pi^{J, [y]} = \{\gamma, u \cdot \beta\}$. Thus Theorem 2.5 says that the order k of the shuttling tour $s_\gamma u s_\beta u^{-1}$ is determined by the equality

$$\langle \gamma, u \cdot \beta \rangle = -\cos(\pi/k) .$$

Moreover, we have $\Phi_J^{\perp [y]} = \Phi_K^{\perp [y] \cap K}$ by definition of K , while $W_J^{\perp [y]}$ is now the dihedral group of order $2k$, having k positive roots. Thus we have another relation

$$|(\Phi_J^{\perp [y]})^+| = |(\Phi_K^{\perp [y] \cap K})^+| = k ,$$

which also enables us to determine the order k if the root system Φ_K of a finite Coxeter group W_K is well understood.

Owing to these arguments, if W_K is not irreducible, namely if t and t' lie in distinct irreducible components of W_J , then we have

$$k = 1 \text{ if } w_y^{t'} \cdot y \neq y, \text{ and } k = 2 \text{ if } w_y^{t'} \cdot y = y .$$

If W_K is irreducible, then the possible situations are completely listed (up to symmetry) in Tables 1 and 2, where we use the canonical labelling $K = \{r_1, \dots, r_n\}$ given in Section 2.2.4, and abbreviate r_i to i .

5 Description of the entire centralizer

Although the subgroup $C_I = C_{x_I, x_I}$ we have described occupies a fairly large part of the centralizer $Z_W(W_I)$, it is in fact not yet the whole of $Z_W(W_I)$. In this section, we describe the structure of the entire centralizer $Z_W(W_I)$ further (Section 5.1). On the other hand, by using the above results, we can also decompose the normalizer $N_W(W_I)$ in a different manner from

Table 1: List of the shuttling tours, in the case Type $K = A_n, B_n$ or D_n

K		(s, s')	(t, t')	order
A_n	$(n \geq 3)$	$(1, 2)$	$(n, n-1)$	1
	$(n = 2)$	$(1, 2)$	$(2, 1)$	3
B_n		$(1, 2)$	$(2, 1)$	4
		$(3, 1)$	$(3, 2)$	2
		$(4, 2)$	$(4, 2)$	2
	$(i \geq 5)$	$(i, i-2)$	$(i, i-2)$	1
	$(i \geq 4)$	$(i, i-1)$	$(i, i-1)$	1
D_n		$(1, 2)$	$(1, 2)$	2
	$(n \geq 6)$	$(4, 2)$	$(4, 2)$	2
	$(4 \neq i \leq n-2)$	$(i, i-2)$	$(i, i-2)$	1
	$(n \geq 5 \text{ even})$	$(n-1, n-2)$	$(n-1, n-2)$	1
	$(n \geq 5 \text{ odd})$	$(n-1, n-2)$	$(n, n-2)$	1

Table 2: List of the shuttling tours, for the remaining cases

K	(s, s')	(t, t')	order	K	(s, s')	(t, t')	order
E_6	$(1, 3)$	$(6, 5)$	1	E_8	$(1, 3)$	$(1, 3)$	1
	$(2, 4)$	$(2, 4)$	3		$(1, 8)$	$(1, 8)$	2
	$(3, 6)$	$(5, 1)$	1		$(2, 4)$	$(2, 4)$	1
E_7	$(1, 3)$	$(1, 3)$	3		$(2, 7)$	$(2, 7)$	1
	$(2, 4)$	$(2, 4)$	1		$(3, 6)$	$(3, 6)$	1
	$(2, 7)$	$(2, 7)$	1		$(7, 1)$	$(7, 1)$	1
	$(3, 6)$	$(3, 6)$	1		$(8, 7)$	$(8, 7)$	3
	$(6, 1)$	$(6, 1)$	2	H_3	$(1, 2)$	$(3, 2)$	2
	$(7, 6)$	$(7, 6)$	1	H_4	$(1, 2)$	$(1, 2)$	3
F_4	$(1, 2)$	$(1, 2)$	3		$(2, 4)$	$(2, 4)$	2
	$(1, 4)$	$(4, 1)$	4		$(4, 3)$	$(4, 3)$	5
	$(2, 4)$	$(3, 1)$	2	$I_2(m)$	$(1, 2)$	$(2, 1)$	m

[5]. We also show the new decomposition of $N_W(W_I)$ as a by-product of our argument (Section 5.2). It is worthy to notice that our decomposition of $N_W(W_I)$ possesses some good properties which failed in [5]; for instance, in contrast with the second factor $\tilde{N}_I$ of the decomposition of $N_W(W_I)$ in [5] which is a Coxeter group as well, our second factor $W^{\perp I}$ is a reflection subgroup of W , not just a general Coxeter group, and the factor admits a simple closed definition.

5.1 On the centralizers

The aim here is to show that $Z_W(W_I)$ is a (not necessarily split) extension of C_I by an elementary abelian 2-group $\tilde{\mathcal{A}}$ described below, and give a further decomposition of $Z_W(W_I)$. Our first observation is that the map $x_\lambda \mapsto y_\lambda$ defines an isomorphism $W_{[x]} \rightarrow W_{[y]}$ for any $x, y \in \mathcal{V}(C)$, since this map is induced by conjugation by an element of $C_{y,x} \neq \emptyset$. Hence if $A \subseteq \Lambda$, the groups W_{x_A} are isomorphic for all $x \in \mathcal{V}(C)$ in this manner, justifying the expression “ A is an irreducible component of Λ ” to mean that W_{x_A} is an irreducible component of $W_{[x]}$. The situation is similar for the expressions “ $A \subseteq \Lambda$ is of finite type” and “ $A \subseteq \Lambda$ is of (-1) -type”. Note that, if $A \subseteq \Lambda$ is of finite type, then

$$ww_0(x_A)w^{-1} = w_0(y_A) \text{ holds in } W \text{ for all } w \in C_{y,x}$$

since the above isomorphism $W_{x_A} \rightarrow W_{y_A}$ sends $w_0(x_A)$ to $w_0(y_A)$.

Let $A \subseteq \Lambda$ be of finite type and a union of irreducible components of Λ . Then define

$$x^A \in S^{(\Lambda)} \text{ for } x \in \mathcal{V}(C) \text{ by } x^A_\lambda = w_0(x_A)x_\lambda w_0(x_A) .$$

Note that there is a common permutation σ_A on Λ such that $x^A_\lambda = x_{\sigma_A(\lambda)}$ for all $x \in \mathcal{V}(C)$ and $\lambda \in \Lambda$, and that $w_0(x_A)$ and $w_0(x_{A'})$ commute (hence σ_A and $\sigma_{A'}$ do as well) for such $A, A' \subseteq \Lambda$. More precisely, $w_0(x_A)w_0(x_{A'}) = w_0(x_{AA'})$ where AA' denotes the symmetric difference of A and A' (hence $AA' = A'A$). Let

$$\tilde{\mathcal{A}} = \{A \subseteq \Lambda \mid x_I^A \text{ is defined and } x_I^A \in \mathcal{V}(C)\} ,$$

which is an elementary abelian 2-group with the symmetric difference as multiplication. Indeed, if $A, A' \in \tilde{\mathcal{A}}$ and $w \in C_{x_I^A, x_I} \neq \emptyset$, then w also lies in $C_{(x_I^A)^{A'}, x_I^{A'}}$ and $x_I^{A'} \in \mathcal{V}(C)$, hence $(x_I^A)^{A'} = x_I^{AA'} \in \mathcal{V}(C)$ and $AA' \in \tilde{\mathcal{A}}$. More precisely, for $A \in \tilde{\mathcal{A}}$, the graph $\mathcal{C}$ defined in Section 3.4 admits an automorphism τ_A such that

$$\begin{aligned} \tau_A(y) &= y^A \text{ for } y \in \mathcal{V}(C), \tau_A(w_y^s) = w_{y^A}^s \text{ for any edge } w_y^s, \\ w_0(z_A)w w_0(y_A) &= \tau_A(w) \text{ holds in } C_{z_A, y^A} \text{ for all } w \in C_{z, y}. \end{aligned} \quad (5.15)$$

Note that the map $A \mapsto \tau_A$ is a group homomorphism $\tilde{\mathcal{A}} \rightarrow \text{Aut } \mathcal{C}$.

Now we have the following result. Recall the maximal tree $\mathcal{T}$ in $\mathcal{Y}^1$ and the paths $p_{y,x}$ in $\mathcal{T}$ introduced in Section 4.3. Note that each path in $\mathcal{C}$ from x to y represents an element of $C_{y,x}$.

Theorem 5.1. *We have an exact sequence $1 \rightarrow C_I \hookrightarrow Z_W(W_I) \xrightarrow{\mathbf{A}} \tilde{\mathcal{A}} \rightarrow 1$ of group homomorphisms and a map $g : \tilde{\mathcal{A}} \rightarrow Z_W(W_I)$ with $\mathbf{A} \circ g = \text{id}_{\tilde{\mathcal{A}}}$, where*

$$\begin{aligned} \mathbf{A} : Z_W(W_I) &\rightarrow \tilde{\mathcal{A}}, w \mapsto \mathbf{A}(w) = \{\lambda \in \Lambda \mid w \cdot \alpha_{(x_I)_\lambda} = -\alpha_{(x_I)_\lambda}\} , \\ g : \tilde{\mathcal{A}} &\rightarrow Z_W(W_I), A \mapsto g_A = p_{x_I, x_I^A} w_0((x_I)_A) . \end{aligned}$$

If we define a subgroup $\tilde{B}_I$ of $Z_W(W_I)$ by

$$\tilde{B}_I = \{w \in Z_W(W_I) \mid (\Phi^{\perp I})^+ = w \cdot (\Phi^{\perp I})^+\} ,$$

then we have $g(\tilde{\mathcal{A}}) \subseteq \tilde{B}_I$, the sequence $1 \rightarrow Y_I \hookrightarrow \tilde{B}_I \xrightarrow{\mathbf{A}} \tilde{\mathcal{A}} \rightarrow 1$ is exact and $Z_W(W_I) = W^{\perp I} \rtimes \tilde{B}_I$, where $\tilde{B}_I$ acts on $W^{\perp I}$ as automorphisms of Coxeter system.

Moreover, if $p \in \pi_1(\mathcal{Y}^1; x_I)$ and $A, A' \in \tilde{\mathcal{A}}$, then we have

$$g_A p g_A^{-1} = \tau_A(p)_{(x_I)} , \quad (5.16)$$

$$g_A g_{A'} = \tau_A(p_{x_I, x_I^{A'}})_{(x_I)} g_{AA'} , \quad (5.17)$$

$$g_A^2 = \tau_A(p_{x_I, x_I^A})_{(x_I)} , \quad (5.18)$$

$$g_A g_{A'} g_A^{-1} g_{A'}^{-1} = \tau_A(p_{x_I, x_I^{A'}})_{(x_I)} \tau_{A'}(p_{x_I^A, x_I})_{(x_I)} . \quad (5.19)$$

Proof. For the first part, the only nontrivial claim is the one on well-definedness of the homomorphism $\mathbf{A}$. This will follow easily once we show that $\mathbf{A}(w) \in \tilde{\mathcal{A}}$. Now Lemma 3.2 implies that $\mathbf{A}(w)$ is of finite type. On the other hand, for $\lambda, \mu \in \Lambda$ with $\langle \alpha_{(x_I)_\lambda}, \alpha_{(x_I)_\mu} \rangle \neq 0$, we have $\lambda \in \mathbf{A}(w)$ if and only if $\mu \in \mathbf{A}(w)$, since w leaves the form $\langle, \rangle$ invariant. Thus $\mathbf{A}(w)$ is a union of irreducible components of Λ , hence $x_I^{\mathbf{A}(w)}$ is defined. Moreover, the remaining claim $x_I^{\mathbf{A}(w)} \in \mathcal{V}(C)$ follows since $w_0((x_I)_{\mathbf{A}(w)})w \in C_{x_I^{\mathbf{A}(w)}, x_I}$. Hence the first part is proven.

For the second part, we have $g_A \in \tilde{B}_I$ for all $A \in \tilde{\mathcal{A}}$, since $p_{x_I, x_I^A} \in Y$ and $w_0((x_I)_A) \in W_I$. This implies the exactness of the sequence. Moreover, the first exact sequence and the properties $g(\tilde{\mathcal{A}}) \subseteq \tilde{B}_I$ and $C_I = W^{\perp I} Y_I$ imply that $Z_W(W_I) = W^{\perp I} \tilde{B}_I$, and then we have $Z_W(W_I) = W^{\perp I} \rtimes \tilde{B}_I$ in the same way as the case of the decomposition $C_I = W^{\perp I} \rtimes Y_I$. The conjugation action of $\tilde{B}_I$ on $W^{\perp I}$ defines automorphisms of Coxeter system since each element of $\tilde{B}_I$ leaves the positive system $(\Phi^{\perp I})^+$ of $W^{\perp I}$ invariant.

Finally, since $w_0(x_{A'}) = w_0(x_A)w_0(x_{AA'}) = w_0((x^{A'})_A)w_0(x_{AA'})$, the last formulae are deduced by straightforward computations based on (5.15). Hence Theorem 5.1 holds. $\square$

For a further study, we define

$$\begin{aligned} \mathcal{A} &= \{A \in \tilde{\mathcal{A}} \mid A \cap A' = \emptyset \text{ if } A' \subseteq \Lambda \text{ is an irreducible component of } (-1)\text{-type}\} , \\ \mathcal{A}' &= \{A \subseteq \Lambda \mid A \text{ is a finite union of irreducible components of } (-1)\text{-type}\} . \end{aligned}$$

Since $x_I^A = x_I$ and $g_A = w_0((x_I)_A)$ for any $A \in \mathcal{A}'$, both $\mathcal{A}$ and $\mathcal{A}'$ are subgroups of $\tilde{\mathcal{A}}$, and we have $\tilde{\mathcal{A}} = \mathcal{A} \times \mathcal{A}'$ and $g(\mathcal{A}') = Z(W_I)$ by the structure of $Z(W_I)$ (see Section 2.2.4). Moreover, $g_{AA'} = g_A g_{A'}$ for $A \in \mathcal{A}$ and $A' \in \mathcal{A}'$. Now the following result is an easy consequence of these properties and Theorem 5.1:

Theorem 5.2. *If we define a subgroup B_I of $\tilde{B}_I$ by*

$$B_I = \{w \in \tilde{B}_I \mid A(w) \in \mathcal{A}\} ,$$

then we have $g(\mathcal{A}) \subseteq B_I$, the sequence $1 \rightarrow Y_I \hookrightarrow B_I \xrightarrow{A} \mathcal{A} \rightarrow 1$ is exact and $\tilde{B}_I = Z(W_I) \times B_I$. Moreover, we have

$$Z_W(W_I) = W^{\perp I} \rtimes (Z(W_I) \times B_I) = (Z(W_I) \times W^{\perp I}) \rtimes B_I ,$$

where B_I acts on $W^{\perp I}$ as automorphisms of Coxeter system.

Note that the exact sequences in these theorems may *not* split, since the map g is *not* necessarily a group homomorphism. However, we have the following result:

Proposition 5.3. *If the maximal tree $\mathcal{T} \subseteq \mathcal{Y}^1$ is stable under the maps τ_A for all $A \in \mathcal{A}$, then $g : \tilde{\mathcal{A}} \rightarrow Z_W(W_I)$ is a group homomorphism. Hence in this case, the exact sequences of Theorems 5.1 and 5.2 split via the g .*

Proof. First, by definition of $\mathcal{A}'$, the τ_A is identity if $A \in \mathcal{A}'$. Then, since $\tilde{\mathcal{A}} = \mathcal{A} \times \mathcal{A}'$, the hypothesis implies that $\mathcal{T}$ is stable under the τ_A for all $A \in \tilde{\mathcal{A}}$. Thus $\tau_A(p_{x_I, x_I^{A'}}) = p_{x_I^A, x_I^{AA'}}$ for all $A, A' \in \tilde{\mathcal{A}}$, hence $g_A g_{A'} = g_{AA'}$ by (5.17) as desired. $\square$

A general result in combinatorial group theory (see e.g., [17, Proposition 10.1]) enables us to construct a presentation of a group extension from those of its factors. In the case of the group B_I , this result gives the following presentation:

Theorem 5.4. *Let $\mathcal{A}_0$ be a basis of the elementary abelian 2-group $\mathcal{A}$. Then B_I admits a presentation, with generators given by $(w_\xi)_{(x_I)}$ for all $w_\xi \in E(\mathcal{Y})$ and g_A for all $A \in \mathcal{A}_0$, and fundamental relations given by those of Y_I , relations (5.16) for $A \in \mathcal{A}_0$ and $q = (w_\xi)_{(x_I)}$ (where $w_\xi \in E(\mathcal{Y})$), relations (5.18) for $A \in \mathcal{A}_0$, and relations (5.19) for $A, A' \in \mathcal{A}_0$.*

Finally, we determine the action of B_I on $W^{\perp I}$ in the following manner:

Proposition 5.5. *Let $w_y^s = s_{\gamma(y,s)}$ be a loop generator of C , where the root $\gamma(x,s)$ is the one defined in Lemma 4.1, and let $w, u \in Y_I$ and $A \in \mathcal{A}$. Then*

$$ur(w, w_y^s)u^{-1} = r(uw, w_y^s) \text{ and } g_A r(w, w_y^s)g_A^{-1} = r(\tau_A(wp_{x_I, y})_{(x_I)}, w_{y^A}^s) .$$

Proof. Since $\tau_A(s_{\gamma(y,s)}) = s_{\gamma(y^A, s)}$, this claim is an easy consequence of (5.15). $\square$

5.2 On the normalizers

In a similar manner, we also give a decomposition of $N_W(W_I)$, in particular that of the second factor N_I of $N_W(W_I) = W_I \rtimes N_I$ (see Section 3.2).

Let $\mathcal{A}_N$ be the set of the bijections $\rho : \Lambda \rightarrow \Lambda$ such that $\rho(x_I) \in \mathcal{V}(C)$, where ρ acts on $S^{(\Lambda)}$ by $\rho(y)_\lambda = y_{\rho^{-1}(\lambda)}$. Then we have $N_I = \bigsqcup_{\rho \in \mathcal{A}_N} C_{x_I, \rho(x_I)}$ by definition. Moreover, the connectedness of $\mathcal{Y}^1$ implies that each ρ fixes all but finitely many elements of Λ , hence ρ has finite order in the symmetric group $\text{Sym}(\Lambda)$ on Λ .

It also holds that each $\rho \in \mathcal{A}_N$ acts on $\mathcal{C}$ as an automorphism by $y \mapsto \rho(y)$ and $w_y^s \mapsto w_{\rho(y)}^s$. In particular, we have $(\rho\rho')(x_I) = \rho(\rho'(x_I)) \in \mathcal{V}(C)$ for all $\rho, \rho' \in \mathcal{A}_N$, where we put $\rho\rho' = \rho \circ \rho'$, therefore $\mathcal{A}_N$ is a subgroup of $\text{Sym}(\Lambda)$. Thus this group $\mathcal{A}_N$ is embedded into $\text{Aut } \mathcal{C}$. Moreover, for a $\rho \in \mathcal{A}_N$ and any path p in $\mathcal{C}$, the two paths p and $\rho(p)$ represent the same element in W .

Now define the following maps

$$\mathbf{A}_N : N_I \rightarrow \mathcal{A}_N, \quad w \mapsto \rho \text{ for any } w \in C_{x_I, \rho(x_I)}$$

and

$$h : \mathcal{A}_N \rightarrow N_I, \quad \rho \mapsto h_\rho = p_{x_I, \rho(x_I)} .$$

Note that $\mathbf{A}_N$ is a group homomorphism since

$$C_{x_I, \rho_1(x_I)} C_{x_I, \rho_2(x_I)} = C_{x_I, \rho_1(x_I)} C_{\rho_1(x_I), \rho_1 \rho_2(x_I)} \subseteq C_{x_I, \rho_1 \rho_2(x_I)} \text{ holds in } W .$$

Then the following analogous results are deduced in a similar way:

Theorem 5.6. *The sequence $1 \rightarrow C_I \hookrightarrow N_I \xrightarrow{\mathbf{A}_N} \mathcal{A}_N \rightarrow 1$ is exact and satisfies that $\mathbf{A}_N \circ h = \text{id}_{\mathcal{A}_N}$. If we define a subgroup $\tilde{Y}_I$ of N_I by*

$$\tilde{Y}_I = \{w \in N_I \mid (\Phi^{\perp I})^+ = w \cdot (\Phi^{\perp I})^+\} ,$$

then $h(\mathcal{A}_N) \subseteq \tilde{Y}_I$, the sequence $1 \rightarrow Y_I \hookrightarrow \tilde{Y}_I \xrightarrow{\mathbf{A}_N} \mathcal{A}_N \rightarrow 1$ is exact and $N_I = W^{\perp I} \rtimes \tilde{Y}_I$, where $\tilde{Y}_I$ acts on $W^{\perp I}$ as automorphisms of Coxeter system. Hence

$$N_W(W_I) = W_I \rtimes (W^{\perp I} \rtimes \tilde{Y}_I) = (W_I \times W^{\perp I}) \rtimes \tilde{Y}_I .$$

Moreover, if $q \in \pi_1(\mathcal{Y}^1; x_I)$, then we have

$$h_\rho^{-1} q h_\rho = \rho^{-1}(q_{(\rho(x_I))}) , \tag{5.20}$$

$$\begin{aligned} h_{\rho_1} h_{\rho_2} h_{\rho_3} \cdots h_{\rho_k} &= p_{x_I, \rho_1(x_I)} \rho_{[1]}(p_{x_I, \rho_2(x_I)}) \rho_{[2]}(p_{x_I, \rho_3(x_I)}) \cdots \\ &\quad \cdots \rho_{[k-1]}(p_{x_I, \rho_k(x_I)}) p_{\rho_{[k]}(x_I), x_I} h_{\rho_{[k]}} , \end{aligned} \tag{5.21}$$

where we put $\rho_{[i]} = \rho_1 \rho_2 \cdots \rho_i$ for each i .

Proposition 5.7. *If the maximal tree $\mathcal{T}$ is stable under all the $\rho \in \mathcal{A}_N$, then h is a group homomorphism. Hence in this case, the exact sequences of Theorem 5.6 split via the h .*

Theorem 5.8 (See [17, Proposition 10.1]). *For the group $\mathcal{A}_N$, choose a generating set $\mathcal{A}'_N$ and fundamental relations of the form “ $\rho_1 \rho_2 \cdots \rho_k = 1$ ” with $\rho_i \in \mathcal{A}'_N$ (this is possible since each $\rho \in \mathcal{A}_N$ has finite order). Then $\tilde{Y}_I$ admits a presentation, with generators given by $(w_\xi)_{(x_I)}$ for all $w_\xi \in E(\mathcal{Y})$ and h_ρ for all $\rho \in \mathcal{A}'_N$, and fundamental relations given by those of Y_I , relations (5.20) for $\rho \in \mathcal{A}'_N$ and $q = (w_\xi)_{(x_I)}$ (where $w_\xi \in E(\mathcal{Y})$), and relations (5.21) for each sequence $\rho_1, \rho_2, \dots, \rho_k \in \mathcal{A}'_N$ such that “ $\rho_1 \rho_2 \cdots \rho_k = 1$ ” is one of the chosen fundamental relations of $\mathcal{A}_N$.*

Proposition 5.9. *Let $w_y^s = s_{\gamma(y,s)}$ be a loop generator of C , and let $w, u \in Y_I$ and $\rho \in \mathcal{A}_N$. Then we have*

$$ur(w, w_y^s) u^{-1} = r(uw, w_y^s) \text{ and } h_\rho r(w, w_y^s) h_\rho^{-1} = r(\rho(wp_{x_I, y})_{(x_I)}, w_{\rho(y)}^s) .$$

6 Example

Let (W, S) be the Coxeter system corresponding to the Coxeter graph in Figure 2, and put $I = \{s_1, s_3, s_4\}$ and $x_I = (s_1, s_3, s_4)$. Then we compute $Z_W(W_I)$ and $N_W(W_I)$.




Figure 2: Coxeter graph for the example (here black circles signify elements of I)

First we construct the graph $\mathcal{C}$. We start with a vertex $x = x_I$, and choose $s \in S \setminus [x]$ such that $[x]_{\sim s}$ is of finite type. Say, $s = s_5$, hence $[x]_{\sim s} = \{s_3, s_4, s_5\}$ and $[x]_{\sim s} \setminus \{s\} = \{s_3, s_4\}$ are of type A_3 and A_2 , respectively; therefore

$$\alpha_{s_3} \xrightarrow{w_0([x]_{\sim s} \setminus \{s\})} -\alpha_{s_4} \xrightarrow{w_0([x]_{\sim s})} \alpha_{s_4} \text{ and } \alpha_{s_4} \xrightarrow{w_0([x]_{\sim s} \setminus \{s\})} -\alpha_{s_3} \xrightarrow{w_0([x]_{\sim s})} \alpha_{s_5} .$$

Thus we have $\varphi(x, s) = ((s_1, s_4, s_5), s_3)$, hence we add to the graph $\mathcal{C}$ a new vertex $y = (s_1, s_4, s_5)$ and a new edge w_x^s from x to y . Iterating such a process, we obtain the entire graph $\mathcal{C}$ finally as depicted in Figure 3 by solid lines. In the figure, we abbreviate s_i to i , and each edge w_y^s is labelled s when the tuples $w_y^s \cdot y$ and y consist of the same contents, since now the element s cannot be determined by the vertices y and $w_y^s \cdot y$ only.




Figure 3: Graph $\mathcal{C}$, complex $\mathcal{Y}$ and shuttling tours for the example

Concerning the circular tours and the shuttling tours, choose a subset $J \subseteq S$ with $|J| = |I| + 2 = 5$, say $J = S \setminus \{s_3\}$, and a connected component $\mathcal{G}$ of $\mathcal{C}(J)$, say the cycle $v_2 v_3 v_4 v_5 v_2$, such that $J_{\sim (J \setminus [y])}$ is of finite type for a vertex y of $\mathcal{G}$. Since $\mathcal{G}$ is now a cycle, it contains a circular tour by definition, yielding a 2-cell of $\mathcal{Y}$ with boundary $\mathcal{G}$. On the other hand, if $J = S \setminus \{s_2\}$,

and $\mathcal{G}$ consists of two loops $w_{v_1}^{s_6}$ and $w_{v_3}^{s_3}$ and a path $p = v_1 v_2 v_3$, then $J_{\sim(J \setminus \{v_1\})} = J \setminus \{s_1\}$ is of finite type, hence we obtain a shuttling tour $w_{v_1}^{s_6} p w_{v_3}^{s_3} p^{-1}$ of order one (see the first row of Table 1, where $K = J \setminus \{s_1\}$). This p is depicted in Figure 3 as a broken line. Iteration of such a process enumerates the circular tours (hence the 2-cells of $\mathcal{Y}$) and the shuttling tours. (Note that the cycle $v_1 v_2 v_5 v_6 v_9 v_{10}$ is *not* the boundary of a 2-cell, since $J_{\sim(J \setminus \{s_1\})} = J$ is *not* of finite type where $J = S \setminus \{s_6\}$.)

We determine the structure of $B_I \subseteq Z_W(W_I)$ and $\widetilde{Y}_I \subseteq N_W(W_I)$. Let e_1, e_2, e_3 be the edges $v_4 \leftarrow v_3, v_6 \leftarrow v_5, v_8 \leftarrow v_7$, respectively, and take a unique maximal tree $\mathcal{T}$ in $\mathcal{Y}^1$ such that $e_1, e_2, e_3 \notin \mathcal{T}$. By Theorems 4.10 and 4.11, $Y_I \simeq \pi_1(\mathcal{Y}; x_I)$ is generated by the three $(e_i)_{(v_1)}$, and has fundamental relations “ $(e_1)_{(v_1)} = 1$ ” and “ $(e_3)_{(v_1)} = 1$ ” induced by the two 2-cells of $\mathcal{Y}$. As a result, Y_I is a free group of rank one (i.e., an infinite cyclic group) generated by $a = (e_2)_{(v_1)}$. Now for B_I , we have $x_I^A = v_{10} \in \mathcal{V}(C)$ and $\mathcal{A} = \{\emptyset, A\}$, where $A = \{2, 3\} \subseteq \Lambda = \{1, 2, 3\}$. Now $\mathcal{T}$ is stable under the map τ_A that sends each v_i to v_{11-i} , therefore Proposition 5.3 implies that B_I is a semidirect product of Y_I by a subgroup $\langle g_A \rangle \simeq \mathcal{A} \simeq \{\pm 1\}$. Moreover, by the formulae in Theorem 5.1, we have

$$g_A a g_A^{-1} = \tau_A(a)_{(v_1)} = e_2^{-1}_{(v_1)} = a^{-1} .$$

Hence, by putting $b = b' = g_A$ and $a' = ba$, we have

$$B_I = \langle a \rangle \rtimes \langle b \rangle \simeq \mathbb{Z} \rtimes \{\pm 1\} \text{ and } B_I = \langle a', b' \mid a'^2 = b'^2 = 1 \rangle \simeq W(\widetilde{A}_1) ,$$

an affine Coxeter group of type $\widetilde{A}_1$.

Similarly, putting $\rho = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix} \in \text{Sym}(\Lambda)$, we have $\mathcal{A}_N = \{\text{id}_\Lambda, \rho\}$, $\rho(v_i) = v_{11-i}$ and $\mathcal{T}$ is stable under the ρ . Thus Theorem 5.6 and Proposition 5.7 imply that

$$h_\rho^{-1} a h_\rho = \rho^{-1}((e_2)_{(v_{10})}) = (e_2)_{(v_1)}^{-1} = a^{-1}$$

and we have $\widetilde{Y}_I = \langle a \rangle \rtimes \langle h_\rho \rangle \simeq W(\widetilde{A}_1)$.

We compute the structure of $W^{\perp I}$ by using Theorems 4.13 and 4.14. Let ξ_i (with $i = 1, 3, 4, 7, 8, 10$) denote the unique pair of the form (v_i, s_j) such that w_{ξ_i} is a loop in $\mathcal{C}$. Then $W^{\perp I}$ is generated by the $r(a^k, w_{\xi_i})$ for all i and $k \in \mathbb{Z}$. For their relations, let $w_{\xi_i} q w_{\xi_j} q^{-1}$ be one of the six shuttling tours. If $(i, j) = (1, 10), (1, 3)$ or $(8, 10)$, then the shuttling tour has order one and $q \subseteq \mathcal{T}$, therefore we have

$$(a^k, w_{\xi_i}) \stackrel{1}{\sim} (a^k q_{(v_1)}, w_{\xi_j}) = (a^k, w_{\xi_j}) .$$

If $(i, j) = (4, 7)$, then the order is also one and $q_{(v_1)} = a^{-1}$, therefore we have

$$(a^k, w_{\xi_4}) \stackrel{1}{\sim} (a^k q_{(v_1)}, w_{\xi_7}) = (a^{k-1}, w_{\xi_7}) .$$

Moreover, if $(i, j) = (3, 4)$ or $(7, 8)$, then the order is two and $q \subseteq \mathcal{T}$, therefore we have

$$(a^k, w_{\xi_i}) \stackrel{2}{\sim} (a^k q_{(v_1)}, w_{\xi_j}) = (a^k, w_{\xi_j}) .$$

These exhaust the relations $\stackrel{m}{\sim}$ on the set $\mathcal{R}_{x_I}$. Thus the generating set R^I of $W^{\perp I}$ consists of infinitely many *distinct* elements

$$r_{1,k} = r(a^k, w_{\xi_1}) \text{ and } r_{4,k} = r(a^k, w_{\xi_4}) , \text{ with } k \in \mathbb{Z}$$

and the fundamental relations are as depicted in Figure 4 where, differently from usual Coxeter graphs, any two adjacent generators commute and a product of any two non-adjacent generators




Figure 4: Structure of the centralizer in the example (here two adjacent generators commute and the product of two non-adjacent generators has infinite order)

has infinite order. In particular, $W^{\perp I}$ is *not* finitely generated though W is finitely generated. Note that this $W^{\perp I}$ possesses no finite irreducible component; this fact can also be deduced directly by using Theorem 4.16.

Finally, we determine the actions of B_I and $\tilde{Y}_I$ on $W^{\perp I}$ by using Propositions 5.5 and 5.9. Put $c = h_\rho$, $\tilde{a} = ca$ and $\tilde{c} = c$. Note that $\{a', b'\}$ and $\{\tilde{a}, \tilde{c}\}$ are the generating sets of B_I and $\tilde{Y}_I$, respectively, as affine Coxeter groups of type $\tilde{A}_1$. Then by the formulae given in those propositions, we have

$$\begin{aligned} a'r_{1,k}a'^{-1} &= \tilde{a}r_{1,k}\tilde{a}^{-1} = r_{1,-k-1} \quad , \quad a'r_{4,k}a'^{-1} = \tilde{a}r_{4,k}\tilde{a}^{-1} = r_{4,-k} \quad , \\ b'r_{1,k}b'^{-1} &= \tilde{c}r_{1,k}\tilde{c}^{-1} = r_{1,-k} \quad , \quad b'r_{4,k}b'^{-1} = \tilde{c}r_{4,k}\tilde{c}^{-1} = r_{4,1-k} \quad . \end{aligned}$$

Thus a' and $\tilde{a}$ act on the graph in Figure 4 as the rotation round the vertical axis through $r_{4,0}$, while b' and $\tilde{c}$ do as the rotation round the vertical axis through $r_{1,0}$. This determines the actions of B_I and $\tilde{Y}_I$.

Finally, we have $Z(W_I) = \langle s_1 \rangle \simeq \{\pm 1\}$ (see Section 2.2.4). Hence the structure of $Z_W(W_I)$ and $N_W(W_I)$ have been completely determined.

7 On the finite part of the factor $W^{\perp I}$

In the final section of this paper, we study further the conjugation action of the group Y_I on the factor $W^{\perp I}$, in particular, on the finite part $W^{\perp I}_{\text{fin}}$ of $W^{\perp I}$. Here $W^{\perp I}_{\text{fin}}$ is defined (as in Section 4.4) to be the product of all the irreducible components of finite type of the Coxeter group $W^{\perp I}$. The result of this section plays an indispensable role in the study on the isomorphism problem in Coxeter groups (see [18]).

We state the main theorem of this section as follows:

Theorem 7.1. *Let $I \subseteq S$. Suppose that I has no irreducible components of type A_n , $2 \leq n < \infty$. Then for any $\gamma \in \Pi^I$, $s_\gamma \in W^{\perp I}_{\text{fin}}$ implies $w \cdot \gamma = \gamma$ for all $w \in Y_I$. Hence every element of the subgroup Y_I of $Z_W(W_I)$ commutes with all elements of $W^{\perp I}_{\text{fin}}$.*

The proof of this theorem will be given below. Before the proof, here we emphasize the following important special case among several cases covered by the theorem:

Corollary 7.2. *Let $I \subseteq S$. If every irreducible component of I of finite type is of (-1) -type (see Section 2.2.4 for the terminology), then we have*

$$Z_W(W_I) = Z(W_I) \times W^{\perp I}_{\text{fin}} \times (W^{\perp I}_{\text{inf}} \rtimes Y_I)$$

where $W^{\perp I}_{\text{inf}}$ denotes the product of all the irreducible components of $W^{\perp I}$ that are not contained in $W^{\perp I}_{\text{fin}}$ (hence $W^{\perp I} = W^{\perp I}_{\text{fin}} \times W^{\perp I}_{\text{inf}}$).

Proof. In this situation, the group $\mathcal{A}$ in Theorem 5.2 is trivial, hence Theorem 5.2 implies that $B_I = Y_I$ and $Z_W(W_I) = Z(W_I) \times (W^{\perp I} \rtimes Y_I)$. Now by Theorem 7.1, both Y_I and $W^{\perp I}_{\text{inf}}$ centralize $W^{\perp I}_{\text{fin}}$, therefore the second factor decomposes as $W^{\perp I}_{\text{fin}} \times (W^{\perp I}_{\text{inf}} \rtimes Y_I)$. Hence Corollary 7.2 holds (assuming Theorem 7.1). $\square$

Regarding the case of a general $I \subseteq S$, the author feels (though the proof has not been given) that the same property would hold for the general case as well. (Note that in the following proof of Theorem 7.1, the assumption on I is not used until the end of the proof of Lemma 7.9.) Thus the author would like to propose the following conjecture:

Conjecture 7.3. *Theorem 7.1 would hold for an arbitrary $I \subseteq S$.*

The rest of this paper is devoted to the proof of Theorem 7.1. First we prepare some temporary terminology. For any $x \in S^*$, we say that a subset Ψ of the simple system $\Pi^{[x]}$ of $W^{\perp [x]}$ is an *irreducible component* of $\Pi^{[x]}$ if $S(\Psi) = \{s_\beta \mid \beta \in \Psi\}$ is an irreducible component of the generating set $R^{[x]}$ of $W^{\perp [x]}$. Such a set Ψ is naturally identified with the vertex set of the Coxeter graph of $(W(\Psi), S(\Psi))$; we will deal with later an automorphism on Ψ in this sense. Note that by Theorem 4.6(3), if $x \in \mathcal{V}(C)$, then the set $\Pi^{[x]}$ consists of the roots $w \cdot \gamma(y, s)$ such that $w \in Y_{x,y}$ and $\gamma(y, s)$ is a positive root defined in Lemma 4.1.

7.1 The first reduction

In the proof of Theorem 7.1, we give several steps to reduce the argument to its special case. The first reduction is given as follows:

Lemma 7.4. *The proof of Theorem 7.1 is reduced to the case that every irreducible component of I is of finite type.*

Proof. Put $x = x_I$. Let $J \subseteq [x]$ denote the union of the irreducible components of $[x]$ that are not of finite type, and let $J' \subseteq S$ denote the union of J and a neighborhood (in the Coxeter graph Γ of W) of each vertex in J . Note that $[x] \cap J' = J$ by the definition of J . Now we show that $Z_W(W_{[x]}) = Z_{W_{S \setminus J'}}(W_{[x] \setminus J'})$ and $\Pi^{[x]} = \Pi^{S \setminus J', [x] \setminus J'}$, which implies the claim of this lemma since every irreducible component of $[x] \setminus J'$ is now of finite type.

By the definitions of J and J' , any generator of the groupoid C of the form w_x^s satisfies that $s \in S \setminus J'$ and $w_x^s \in W_{S \setminus J'}$, and we have $[y] \cap J' = J = [x] \cap J'$ where $y = w_x^s \cdot x$. Iterating this argument, it follows that any generator w_y^s of C satisfies that $s \in S \setminus J'$, $w_y^s \in W_{S \setminus J'}$ and $[y] \cap J' = J$. Moreover, we have $x_A \subseteq S \setminus J'$ for any $A \in \tilde{\mathcal{A}}$ (see Section 5.1 for the notation). This implies that both C_I and $A(\tilde{\mathcal{A}})$ are contained in $W_{S \setminus J'}$ (see Theorem 5.1 for the notation), and we have $Z_W(W_I) \subseteq W_{S \setminus J'}$ since $Z_W(W_I)$ is generated by C_I and $A(\tilde{\mathcal{A}})$ by Theorem 5.1. Hence we have $Z_W(W_{[x]}) = Z_{W_{S \setminus J'}}(W_{[x] \setminus J'})$ by the definition of J' . Moreover, the above argument and the above description of elements of $\Pi^{[x]}$ imply that $\Pi^{[x]} \subseteq \Phi_{S \setminus J'}^{\perp [x] \setminus J'}$, therefore $\Phi^{\perp [x]} = \Phi_{S \setminus J'}^{\perp [x] \setminus J'}$ and $\Pi^{[x]} = \Pi^{S \setminus J', [x] \setminus J'}$. Hence Lemma 7.4 holds. $\square$

7.2 The second reduction

By virtue of Lemma 7.4, we may assume without loss of generality that every irreducible component of I is of finite type. Let $w \in Y_I$ and $\gamma \in \Pi^I$, and suppose that $s_\gamma \in W^{\perp I}_{\text{fin}}$. Let Ψ denote the union of the irreducible components of Π^I containing $w^k \cdot \gamma$ for some $k \in \mathbb{Z}$. Now we give the second reduction of the proof to its special case as follows:

Lemma 7.5. *The proof of Theorem 7.1 is reduced further to the following case: There exists a subset $J \subseteq S$ of finite type such that $I \setminus J$ and J are not adjacent in the Coxeter graph Γ of W and $\Psi \cup \Pi_{I \cap J}$ is a basis of V_J .*

Proof. Put $x = x_I$. First, we show that $|\Psi| < \infty$ and Ψ is of finite type. Take a finite subset K of S such that $w \in W_K$ and $\gamma \in \Phi_K$. Then the number of mutually orthogonal roots of the form $w^k \cdot \gamma$ is at most $|K|$, since those roots are linearly independent and contained in the $|K|$ -dimensional vector space V_K . This implies that the number of irreducible components of $\Pi^{[x]}$ containing some $w^k \cdot \gamma$, which are of finite type by the property $s_\gamma \in W^{\perp [x]}_{\text{fin}}$ and Theorem 4.6(4), is finite, therefore the union Ψ of those irreducible components is also of finite type, in particular of finite cardinality.

Secondly, we show that $[x] \setminus L$ and $\text{Supp } \Psi$ are not adjacent in Γ , where $L = [x] \cap \text{Supp } \Psi$. Assume contrary that $x_i \in [x] \setminus L$ and $s \in \text{Supp } \Psi$ are adjacent in Γ . By the definition of $\text{Supp } \Psi$, we have $s \in \text{Supp } \beta \subset \text{Supp } \Psi$ for some $\beta \in \Psi$. Let $c > 0$ be the coefficient of α_s in β . Then the property $x_i \notin L$ implies that $x_i \notin \text{Supp } \beta$, therefore $\langle \alpha_{x_i}, \beta \rangle \leq c \langle \alpha_{x_i}, \alpha_s \rangle < 0$. This contradicts the property $\beta \in \Phi^{\perp [x]}$. Hence the claim of this paragraph holds.

Put $\Psi' = \Psi \cup \Pi_L$. Then we have $\text{Supp } \Psi' = \text{Supp } \Psi$, therefore $[x] \setminus \text{Supp } \Psi'$ and $\text{Supp } \Psi'$ are not adjacent in Γ . On the other hand, since $|\text{Supp } \Psi'| < \infty$ (recall that $|\Psi| < \infty$) and every irreducible component of $[x]$ is of finite type, it follows that Π_L is of finite type as well as Ψ . Since $\Psi \subset \Pi^{\perp [x]}$, Ψ' is a root basis (see Theorem 2.5) with $|W(\Psi')| < \infty$, hence Ψ' is a basis of a subspace $U = \text{span } \Psi' \subseteq V_{\text{Supp } \Psi'}$ by Corollary 2.6. By applying Proposition 2.7 to $W_{\text{Supp } \Psi'}$ instead of W , there exist $u \in W_{\text{Supp } \Psi'}$ and $J \subseteq \text{Supp } \Psi'$ such that W_J is finite, $u \cdot (U \cap \Phi^+) = \Phi_J^+$ and $u \cdot (U \cap \Pi) \subseteq \Pi_J$. Note that $u \cdot \Psi'$ is a basis of V_J since Ψ' is a basis of U . Now we have $u \cdot \Pi_L \subseteq \Pi_J$ since $\Pi_L \subseteq U \cap \Pi$, while u fixes $\Pi_{[x] \setminus L}$ pointwise since $[x] \setminus \text{Supp } \Psi'$ and $\text{Supp } \Psi'$ are not adjacent in Γ . This implies that $u \in C_{y,x}$ for some $y \in \mathcal{V}(C)$ such that $[y] \cap \text{Supp } \Psi' \subseteq J$ and $[y] \setminus \text{Supp } \Psi' = [x] \setminus \text{Supp } \Psi'$, hence $[y] \cap J$ and J are not adjacent in Γ .

Finally, suppose that $u \cdot \beta' \in \Phi^-$ for some $\beta' \in \Pi^{[x]} \setminus \Psi$. Then we have $\beta' \in \Phi_{\text{Supp } \Psi'}$ and $s_{\beta'} \in W_{\text{Supp } \Psi'}$, while β' is orthogonal to Ψ' by the definition of Ψ' . This implies that the element $us_{\beta'}$ also satisfies the above characteristics of the element u , and we have $\ell(us_{\beta'}) < \ell(u)$ by the choice of β' . Hence by choosing such an element u of minimal length, it follows that $u \cdot \beta' \in \Phi^+$ for every $\beta' \in \Pi^{[x]} \setminus \Psi$, therefore $u \cdot \Pi^{[x]} \subseteq \Phi^+$ and $u \in Y_{y,x}$. Since $uwu^{-1} \in Y_{y,y}$ fixes $u \cdot \gamma \in \Pi^{[y]}$ if and only if w fixes γ , the claim of Lemma 7.5 holds by letting $y \in \mathcal{V}(C)$, $uwu^{-1} \in Y_{y,y}$, $u \cdot \gamma \in \Pi^{[y]}$ and $u \cdot \Psi \subseteq \Pi^{[y]}$ play the roles of $x = x_I$, w , γ and Ψ , respectively (see also Theorem 4.6(4)). $\square$

From now on, we assume that the condition in the statement of Lemma 7.5 is satisfied. Moreover, we may assume further that each irreducible component J' of J is not contained in I ; indeed, if $J' \subseteq I$, then we can replace J with the smaller set $J \setminus J'$ that also satisfies the characteristics of the set J in Lemma 7.5. Now there exist two possibilities; either $\Psi \not\subseteq \Phi_{I^\perp}$, or $\Psi \subseteq \Phi_{I^\perp}$ (see (4.14) for the notation $I^\perp$). We discuss the first case in Section 7.4 and the second case in Section 7.5.

7.3 Another decomposition for elements of Y

Before starting the discussion of those two cases, we introduce another decomposition of an element of $u \in Y$ in the following inductive manner, together with some relevant notations and terminology. Let $u \in Y_{z,y}$, and put $u^{(0)} = u$ and $y^{(0)} = y$. Let $J^{(0)} \subseteq S$ such that $u \cdot \Pi_{J^{(0)}} \subseteq \Pi$. Now suppose that $i \geq 0$ and we have already defined $J^{(i)}$, $y^{(i)}$ and $u^{(i)} \in Y_{z,y^{(i)}}$ with the property $u^{(i)} \cdot \Pi_{J^{(i)}} \subseteq \Pi$. If $u^{(i)} = 1$ then the task is over. Suppose $u^{(i)} \neq 1$, and take a $t^{(i)} \in S$ such that $u \cdot \alpha_{t^{(i)}} \in \Phi^-$. Note that $t^{(i)} \notin [y^{(i)}] \cup J^{(i)}$ and $t^{(i)}$ is adjacent to $[y^{(i)}]$ in Γ since $u^{(i)} \in Y_{z,y^{(i)}}$. Put $K = ([y^{(i)}] \cup J^{(i)})_{\sim t^{(i)}} \subseteq S$. Then by Lemma 3.2, K is of finite type and the element

$$\omega = \omega_{y^{(i)}, J^{(i)}}^{t^{(i)}} = w_0(K)w_0(K \setminus \{t^{(i)}\})$$

is a right divisor of $u^{(i)}$. We call the set K the *support* of $\omega_{y^{(i)}, J^{(i)}}^{t^{(i)}}$. By the definition, there exist unique $y^{(i+1)} \in \mathcal{V}(C)$ and $J^{(i+1)} \subseteq S$ such that $\omega \cdot y^{(i)} = y^{(i+1)}$ and $\omega \cdot \Pi_{J^{(i)}} = \Pi_{J^{(i+1)}}$. Moreover, Lemma 2.2 implies that $\Phi[\omega] \subseteq \Phi[u^{(i)}]$, therefore $\Phi^\perp[y^{(i)}][\omega] = \emptyset$ and $\omega \in Y_{y^{(i+1)}, y^{(i)}}$. Thus we have $u^{(i+1)} = u^{(i)}\omega^{-1} \in Y_{z, y^{(i+1)}}$, $u^{(i+1)} \cdot \Pi_{J^{(i+1)}} \subseteq \Pi$ and we obtain (inductively) a partial decomposition

$$u = u^{(i+1)}\omega_{y^{(i)}, J^{(i)}}^{t^{(i)}}\omega_{y^{(i-1)}, J^{(i-1)}}^{t^{(i-1)}} \cdots \omega_{y^{(1)}, J^{(1)}}^{t^{(1)}}\omega_{y^{(0)}, J^{(0)}}^{t^{(0)}}$$

of u , which becomes the desired decomposition when $u^{(i+1)} = 1$. We call each $\omega_{y^{(i)}, J^{(i)}}^{t^{(i)}}$ a *wide transformation* if its support K contains an element of $J^{(i)} \setminus [y^{(i)}]$ and a *narrow transformation* if K contains no elements of $J^{(i)} \setminus [y^{(i)}]$ (in the latter case $\omega_{y^{(i)}, J^{(i)}}^{t^{(i)}}$ coincides with a non-loop generator $w_{y^{(i)}}^{t^{(i)}}$ of Y). Decompositions for elements of Y of the above type will be used in the following argument.

Here we notice some properties of the above decomposition as follows. First, by the definition, we have $\ell(u) = \sum_{j=0}^n \ell(\omega_{y^{(j)}, J^{(j)}}^{t^{(j)}})$ for any decomposition $u = \omega_{y^{(n)}, J^{(n)}}^{t^{(n)}} \cdots \omega_{y^{(1)}, J^{(1)}}^{t^{(1)}}\omega_{y^{(0)}, J^{(0)}}^{t^{(0)}}$ of $u \in Y_{z,y}$ as above. Secondly, we have the following:

Lemma 7.6. *Let $u = \omega_{y^{(n)}, J^{(n)}}^{t^{(n)}} \cdots \omega_{y^{(1)}, J^{(1)}}^{t^{(1)}}\omega_{y^{(0)}, J^{(0)}}^{t^{(0)}}$ be a decomposition of $u \in Y_{z,y}$ as above. Then for each $0 \leq i \leq n$, there exists an element of $\Pi_{K \setminus \{t^{(i)}\}}$, where K is the support of $\omega = \omega_{y^{(i)}, J^{(i)}}^{t^{(i)}}$, which is not fixed by ω .*

Proof. Assume contrary that ω fixes $\Pi_{K \setminus \{t^{(i)}\}}$ pointwise. Then by the same argument as the one deriving the second condition in Lemma 4.1 from the first condition in that lemma, there exists a root $\gamma \in (\Phi_K^{\perp K \setminus \{t^{(i)}\}})^+$ such that $\omega \cdot \gamma \in \Phi^-$. Now this γ also belongs to $(\Phi^\perp[y^{(i)}])^+$, therefore $\Phi^\perp[y^{(i)}][\omega] \neq \emptyset$, while ω is a right divisor of $\omega_{y^{(n)}, J^{(n)}}^{t^{(n)}} \cdots \omega_{y^{(i)}, J^{(i)}}^{t^{(i)}} \in Y_{z, y^{(i)}}$ by the above property. This contradicts Lemma 2.2 and the definition of Y . Hence Lemma 7.6 holds. $\square$

On the other hand, the following lemma is straightforward:

Lemma 7.7. *Let $u = \omega_n \cdots \omega_1 \omega_0$, where $\omega_i = \omega_{y^{(i)}, J^{(i)}}^{t^{(i)}}$ for each $0 \leq i \leq n$, be a decomposition of $u \in Y_{z,y}$ as above. Let $0 \leq i_1 < i_2 < \cdots < i_k \leq n$ be indices such that for each j , the action of ω_{i_j} leaves the sets $[y^{(i_j)}]$ and $\Pi_{J^{(i_j)}}$ invariant. Let $\hat{u} = \omega_n \cdots \hat{\omega}_{i_k} \cdots \hat{\omega}_{i_1} \cdots \hat{\omega}_{i_0} \cdots \omega_0$ be the product obtained from $\omega_n \cdots \omega_1 \omega_0$ by removing every ω_{i_j} , $1 \leq j \leq k$. Let σ denote the map from $\{0, 1, \dots, n-k\}$ to $\{0, 1, \dots, n\}$ such that the j -th factor (from right) in $\hat{u}$ is $\omega_{\sigma(j)}$ for every $0 \leq j \leq n-k$. Then there exist $y'^{(j)} \in \mathcal{V}(C)$ for $0 \leq j \leq n-k+1$ such that $y'^{(0)} = y^{(0)}$, $[y'^{(j+1)}] = [y^{(\sigma(j)+1)}] = [y^{(\sigma(j+1))}]$ and $\omega_{\sigma(j)} \in Y_{y'^{(j+1)}, y'^{(j)}}$ for every $0 \leq j \leq n-k$, where $y^{(n+1)} = z$. Hence $\hat{u} \in Y_{y'^{(n+1)}, y'}$.*

Moreover, we have the following lemma:

Lemma 7.8. *Let $u = \omega_n \cdots \omega_1 \omega_0$ be a product of some elements $\omega_i = \omega_{y^{(i)}, J^{(i)}}^{t^{(i)}} \in Y_{y^{(i+1)}, y^{(i)}}$ as above such that $\omega_i \cdot \Pi_{J^{(i)}} = \Pi_{J^{(i+1)}}$ for every $0 \leq i \leq n$, hence $u \in Y_{y^{(n+1)}, y^{(0)}}$. Here we do not assume that the length of u is the sum of those of the ω_i . Put $y = y^{(0)}$ and $z = y^{(n+1)}$. Suppose that there exists an index $\lambda \in \Lambda$ such that the support of every ω_i does not contain y_λ . Let $s \in J^{(0)}$ and $s' \in J^{(n+1)}$ such that $u \cdot \alpha_s = \alpha_{s'}$. Then we have $z_\lambda = y_\lambda$ and $u \in \hat{Y}_{z', y'}$ (see Section 3.1 for the definition of $\hat{Y}$), where y' and z' are elements of $S^{(\Lambda)}$ obtained from y and z by replacing $y_\lambda = z_\lambda$ with s and with s' , respectively.*

Proof. First, since the support K_0 of ω_0 does not contain y_λ , it follows that K_0 is not adjacent to y_λ and $(y^{(1)})_\lambda = y_\lambda$. Secondly, since $\omega_0 \cdot \Pi_{J^{(0)}} = \Pi_{J^{(1)}}$, there is an $s_1 \in J^{(1)}$ such that $\omega_0 \cdot \alpha_s = \alpha_{s_1}$. Let $y'^{(1)}$ be the element of $S^{(\Lambda)}$ obtained from $y^{(1)}$ by replacing $y_\lambda = (y^{(1)})_\lambda$ with s_1 . Then we have $\omega_0 \cdot y' = y'^{(1)}$ by the construction. Moreover, by the definition of ω_0 , the sets $([y] \cup J^{(0)}) \setminus K_0$ and K_0 are not adjacent to each other, therefore $([y'] \cup J^{(0)}) \setminus K_0$ (which is a subset of $([y] \cup J^{(0)}) \setminus K_0$) and K_0 are not adjacent to each other. Since $[y] \cap K_0 \subseteq [y'] \cap K_0$, it follows that $\Phi^{\perp[y']}[\omega_0] = \Phi_{K_0}^{\perp[y'] \cap K_0}[\omega_0] \subseteq \Phi_{K_0}^{\perp[y] \cap K_0}[\omega_0] = \Phi^{\perp[y]}[\omega_0] = \emptyset$, therefore we have $\omega_0 \in \hat{Y}_{y'^{(1)}, y'}$. Now the claim follows by iterating the argument inductively. $\square$

7.4 The first case $\Psi \not\subseteq \Phi_{I^\perp}$

In this subsection, we consider the case that $\Psi \not\subseteq \Phi_{I^\perp}$ among the two cases mentioned in the last of Section 7.2. Note that if w fixes Ψ pointwise then the claim follows, since Ψ contains γ by the definition. First we present an important lemma for the analysis of this case as follows:

Lemma 7.9. *In the situation, the subset J of S is irreducible.*

Proof. The condition $\Psi \not\subseteq \Phi_{I^\perp}$ and the characteristics of J imply that there exists an irreducible component K of J that contains an element of I (and is not contained in I by our assumption). Since the support of any root is irreducible, by putting $\Psi' = \Psi \cap \Phi_K$ we have $\Psi' \neq \emptyset$ and $\Psi' \cup \Pi_{I \cap K}$ is a basis of V_K (recall that $\Psi \cup \Pi_{I \cap J}$ is a basis of V_J). Note that Ψ' is a union of irreducible components of Ψ since Ψ' is orthogonal to $\Psi \setminus \Psi'$ by the definition of Ψ' , and that $\langle w \rangle \subseteq W$ acts transitively on the finite set of irreducible components of Ψ by the definition of Ψ . Now the claim of Lemma 7.9 follows if we have $w \cdot \Phi_K = \Phi_K$. Indeed, if $w \cdot \Phi_K = \Phi_K$, then we have $w \cdot \Psi' = \Psi'$, therefore $\Psi = \Psi'$ by the transitivity. Hence we have $\Psi \subseteq \Phi_K$ and $J = K$ is irreducible as desired, since we have assumed that each irreducible component of J is not contained in I (see the final paragraph of Section 7.2).

From now, we show that $w \cdot \Phi_K = \Phi_K$. For the purpose, it suffices to show that $w \cdot \Phi_K \subseteq \Phi_K$ (since K is of finite type as well as J), or equivalently $w \cdot \Pi_K \subseteq \Phi_K$. Moreover, since K is irreducible, $K \cap I \neq \emptyset$ and $w \cdot \Pi_{K \cap I} = \Pi_{K \cap I}$, it suffices to show that $w \cdot \alpha_s \in \Phi_K$ provided s is adjacent to some $t \in K$ in Γ such that $w \cdot \alpha_t \in \Phi_K$. Now note that $w \cdot \Psi = \Psi$ and $w \cdot \Pi_{I \cap J} = \Pi_{I \cap J}$, therefore $w \cdot \Phi_J = \Phi_J$. Assume contrary that $w \cdot \alpha_s \notin \Phi_K$. Then we have $w \cdot \alpha_s \in \Phi_J \setminus \Phi_K = \Phi_{J \setminus K}$ since K is an irreducible component of J , therefore $w \cdot \alpha_s$ is orthogonal to $w \cdot \alpha_t \in \Phi_K$. This contradicts the property that s is adjacent to t , since w leaves the form $\langle \cdot, \cdot \rangle$ invariant. Hence we have $w \cdot \alpha_s \in \Phi_K$ as desired, therefore Lemma 7.9 holds. $\square$

By virtue of Lemma 7.9, we can apply the classification of finite irreducible Coxeter groups to the subset J of S . We also use the concrete descriptions of root systems of finite irreducible Coxeter groups (see e.g., [16, Sections 2.10 and 2.13]). Let $J = \{r_1, r_2, \dots, r_N\}$ be the canonical labelling given in Section 2.2.4. Write $\alpha_i = \alpha_{r_i}$ for simplicity. Let $x = x_I$, and write $x =$

$(x_{(J)}; x_{(J^c)})$ where $x_{(J)} \in S^*$ consists of the elements of $[x] \cap J$ and $x_{(J^c)} \in S^*$ denotes the remaining part of x .

Before starting case-by-case analysis based on the classification of J , we give a remark to simplify the argument. Suppose that we have given a collection of $y_{(J)} \in S^*$ such that $[y_{(J)}] \subseteq J$ and for any possible choice of x , there exists an element $u \in Y \cap W_J$ such that $u \cdot x_{(J)} = y_{(J)}$ (hence $y = u \cdot x = (y_{(J)}; x_{(J^c)})$). Then by letting y , uwu^{-1} , and $u \cdot \Psi$ play the roles of x , w , and Ψ , respectively, we may assume without loss of generality that $x_{(J)}$ coincides with one of these $y_{(J)}$. Thus the width of our case-by-case analysis can be reduced by focusing on such “representatives” $y_{(J)}$. Moreover, note that now $\Psi = \Pi^{J, I \cap J}$, since otherwise the subset $\Psi \cup \Pi_{I \cap J}$ of $\Pi^{J, I \cap J} \cup \Pi_{I \cap J} \subseteq V_J$ cannot span V_J .

Now we divide the following argument according to the type of J , where our aim is to show that w fixes Ψ pointwise (which implies our claim as mentioned above):

Type A_N ($N \geq 2$) First, note that the description of actions of the longest elements of finite parabolic subgroups given in Section 2.2.4 implies the following two properties:

- If $z = (r_k)$ and $s = r_{k+1}$, then $w_z^s \cdot z = (r_{k+1})$.
- If $z = (r_k, r_{k+2})$ and $s = r_{k+1}$, then $w_z^s \cdot z = (r_{k+2}, r_k)$.

By using these two kinds of transformations, we can choose above-mentioned representatives $y_{(J)}$ each of which consists of the elements $r_N, r_{N-2}, \dots, r_{N-2k}$, where $k \in \mathbb{Z}$ and $0 \leq k \leq (N-1)/2$ (recall that I is assumed to have no irreducible components of type A_n , $n \geq 2$). If $x_{(J)}$ is the above member of representatives, then by the shape of the root system of type A_N , the set $\Psi = \Pi^{J, [x_{(J)}]}$ consists of α_j with $1 \leq j \leq N-2k-2$. This implies that $\Psi \cup \Pi_{[x_{(J)}]}$ cannot span V_J , contradicting the current situation. Hence in fact this case does not occur.

Type B_N ($N \geq 2$) By a similar argument, we may assume without loss of generality that $x_{(J)}$ satisfies one of the following two conditions:

- $x_{(J)}$ consists of $r_N, r_{N-1}, \dots, r_{N-l}, r_{N-l-2}, r_{N-l-4}, \dots, r_{N-l-2k}$, where $k, l \in \mathbb{Z}$, $k \geq 0$, $l \geq 0$ and $l+2k \leq N-1$.
- $x_{(J)}$ consists of $r_{N-1}, r_{N-3}, \dots, r_{N-2k-1}$, where $k \in \mathbb{Z}$ and $0 \leq k \leq (N-2)/2$.

In the first case, put $m = N-l-2k$. Then $\Psi = \Pi^{J, [x_{(J)}]}$ consists of α_j for $1 \leq j \leq m-2$, $\beta = \sum_{j=m-1}^{N-1} \sqrt{2}\alpha_j + \alpha_N$ (when $m \geq 2$), and $\beta'_i = \alpha_{m+2i} + \sum_{j=m+2i+1}^{N-1} 2\alpha_j + \sqrt{2}\alpha_N$ for $0 \leq i \leq k-1$. Now if $k \geq 1$, then the facts $w \cdot \beta'_{k-1} \in \Psi$ (since $w \cdot \Psi = \Psi$) and

$$w \cdot \alpha_{N-l-1} = (w \cdot \beta'_{k-1} - \alpha_{N-l-2} - \sum_{j=N-l}^{N-1} 2\alpha_j - \sqrt{2}\alpha_N)/2 \in \Phi$$

imply that $w \cdot \beta'_{k-1} = \beta'_{k-1}$, while $\{\beta'_{k-1}\}$ is an irreducible component of Ψ , therefore we have $\Psi = \{\beta'_{k-1}\}$ (since $\langle w \rangle$ acts transitively on the set of the irreducible components of Ψ by the definition of Ψ) and the claim follows.

On the other hand, if $k = 0$, then $\Psi = \{\alpha_1, \dots, \alpha_{N-l-2}, \beta\}$ is of type B_{N-l-1} with the ordering of elements compatible to the canonical labelling in Section 2.2.4, where we put $B_1 = A_1$ (note that now $N-l \geq 2$ since $\Psi \neq \emptyset$). Now the facts $w \cdot \beta \in \Psi$ and

$$w \cdot \alpha_{N-l-1} = (w \cdot \beta - \sum_{j=N-l}^{N-1} \sqrt{2}\alpha_j - \alpha_N)/\sqrt{2} \in \Phi$$

imply that $w \cdot \beta = \beta$, therefore (even if $N - l = 3$ and Ψ is of type B_2) the automorphism on Ψ induced by the action of w is trivial. Thus w fixes Ψ pointwise, as desired.

In the second case, put $m = N - 2k - 1$. Then $\Psi = \Pi^{J, [x_{(J)}]}$ consists of α_j for $1 \leq j \leq m - 2$, $\beta = \sum_{j=m-1}^{N-1} \sqrt{2}\alpha_j + \alpha_N$ (when $m \geq 2$), and $\beta'_i = \alpha_{m+2i} + \sum_{j=m+2i+1}^{N-1} 2\alpha_j + \sqrt{2}\alpha_N$ for $0 \leq i \leq k$. Now the facts $w \cdot \beta'_k \in \Psi$ and $w \cdot \alpha_N = (w \cdot \beta'_k - \alpha_{N-1})/\sqrt{2} \in \Phi$ imply that $w \cdot \beta'_k = \beta'_k$, while $\{\beta'_k\}$ is an irreducible component of Ψ , therefore we have $\Psi = \{\beta'_k\}$ and the claim follows. Hence the claim holds in any case of type B_N .

Type D_N ($N \geq 4$) By a similar argument and symmetry, we may assume without loss of generality that $x_{(J)}$ satisfies one of the following two conditions:

- $x_{(J)}$ consists of $r_N, r_{N-1}, \dots, r_{N-l}, r_{N-l-2}, \dots, r_{N-l-2k}$, where $k, l \in \mathbb{Z}$, $l \geq 1$, $k \geq 0$, and $l + 2k \leq N - 1$.
- $x_{(J)}$ consists of $r_{N-1}, r_{N-3}, \dots, r_{N-2k-1}$, where $k \in \mathbb{Z}$ and $0 \leq k \leq (N - 2)/2$.

We consider the first choice of $x_{(J)}$. Put $m = N - l - 2k$. Now the set $\Psi = \Pi^{J, [x_{(J)}]}$ consists of α_j for $1 \leq j \leq m - 2$, and $\beta_i = \alpha_{m-2+2i} + \sum_{j=m-1+2i}^{N-2} 2\alpha_j + \alpha_{N-1} + \alpha_N$ for $0 \leq i \leq k$ (where β_0 is defined only when $m \geq 3$). Since $\Psi \cup \Pi_{[x_{(J)}]}$ spans V_J , it is necessary that $m \neq 2$. If $k \geq 1$, then the facts $w \cdot \beta_k \in \Psi$ and

$$w \cdot \alpha_{N-l-1} = (w \cdot \beta_k - \alpha_{N-l-2} - \sum_{j=N-l}^{N-2} 2\alpha_j - \alpha_{N-1} - \alpha_N)/2 \in \Phi$$

imply that $w \cdot \beta_k = \beta_k$, while $\{\beta_k\}$ is an irreducible component of Ψ . Since $\langle w \rangle$ acts transitively on the set of irreducible components of Ψ , it follows that $\Psi = \{\beta_k\}$ and the claim holds.

On the other hand, assume $k = 0$. Since $\Psi \neq \emptyset$, we have $m = N - l \geq 3$ and $\Psi = \{\alpha_1, \alpha_2, \dots, \alpha_{m-2}, \beta_0\}$ is of type D_{m-1} , where we put $D_2 = A_1 \times A_1$ and $D_3 = A_3$. Now we give the following lemma:

Lemma 7.10. *In this setting, we have $\{w \cdot \alpha_{m-2}, w \cdot \beta_0\} = \{\alpha_{m-2}, \beta_0\}$ and $w \cdot \alpha_i = \alpha_i$ for all $1 \leq i \leq m - 3$.*

Proof. First, note that

$$w \cdot \alpha_{m-1} = (w \cdot \beta_0 - w \cdot \alpha_{m-2} - \sum_{j=m}^{N-2} 2\alpha_j - \alpha_{N-1} - \alpha_N)/2 \in \Phi.$$

This implies that for every $1 \leq j \leq m - 2$, the coefficients of α_j in $w \cdot \beta_0$ and in $w \cdot \alpha_{m-2}$ (both being integers) have the same parity. By the above description of Ψ , this is possible only if w leaves the set $\{\beta_0, \alpha_{m-2}\}$ invariant. Moreover, this implies that the automorphism on Ψ induced by the action of w must fix every α_j , $1 \leq j \leq m - 3$. Hence Lemma 7.10 holds. $\square$

If $w \cdot \beta_0 = \beta_0$, then w fixes Ψ pointwise as desired. Assume contrary that $w \cdot \beta_0 = \alpha_{m-2}$ and $w \cdot \alpha_{m-2} = \beta_0$ to deduce a contradiction. Then we have $w \cdot \alpha_{m-1} \in \Phi_J^-$. By Lemma 3.2, w_x^s is a right divisor of w , where $s = r_{m-1}$. Moreover, the action of w_x^s on x exchanges r_{N-1} and r_N and fixes the remaining points (note that $[x]_{\sim s} = [x_{(J)}] \cup \{s\}$), therefore $w_x^s \in Y$ by Lemma 4.1. Put $y = w_x^s \cdot x$ and $u = w(w_x^s)^{-1}$. Then we have $u \in Y_{x,y}$ and u fixes Ψ pointwise, therefore we have $u \cdot \alpha_{m-1} = \alpha_{m-1}$ and $u \cdot \Pi_J = \Pi_J$. Now consider a decomposition of u as in Section 7.3 with respect to $J^{(0)} = J$. Note that the action of u does not fix $r_N \in [y] \cap J$ and J is irreducible. This implies that the decomposition of u contains at least one wide transformation.

Let $\omega = \omega_{y^{(i)}, J^{(i)}}^{t^{(i)}}$ denote the wide transformation with least index $i \geq 0$. Then the action of the element $u' = \omega_{y^{(i-1)}, J^{(i-1)}}^{t^{(i)}} \cdots \omega_{y^{(0)}, J^{(0)}}^{t^{(0)}}$ fixes every element of Π_J . Since the support K of ω is irreducible and of finite type, $t^{(i)}$ is adjacent to $[y^{(i)}]$, and I (hence $[y^{(i)}]$) has no irreducible components of type A_n with $n \geq 2$, we have the following seven possibilities (up to symmetry) for the choice of $t = t^{(i)}$ and K :

1. $K = J \cup \{t, t'\}$ is of type D_{N+2} , $t' \in [y^{(i)}]$, and $m_{t, r_1} = m_{t', t} = 3$.
2. $N = 7$, $K = J \cup \{t\}$ is of type E_8 , and $m_{t, r_6} = 3$.
3. $N = 6$, $K = J \cup \{t\}$ is of type E_7 , and $m_{t, r_5} = 3$.
4. $N = 5$, $K = J \cup \{t, t'\}$ is of type E_7 , $t' \in [y^{(i)}]$, and $m_{t, r_4} = m_{t', t} = 3$.
5. $N = 5$, $K = J \cup \{t\}$ is of type E_6 , and $m_{t, r_4} = 3$.
6. $N = 4$, $K = J \cup \{t, t'\}$ is of type D_6 , $t' \in [y^{(i)}]$, and $m_{t, r_3} = m_{t', t} = 3$.
7. $N = 4$, $K = J \cup \{t\}$ is of type D_5 , and $m_{t, r_3} = 3$.

We show that the cases 1, 3, and 6 do not occur. Namely, in these cases ω fixes every point in the set $\Pi_{K \setminus \{t\}}$, contradicting Lemma 7.6. Hence these cases do not occur, as desired.

We show that the cases 2 and 4 do not occur. Put

$$\gamma = \begin{cases} \alpha_2 + 2\alpha_3 + 3\alpha_4 + 4\alpha_5 + 3\alpha_6 + 2\alpha_7 + 2\alpha_t & \text{in the case 2,} \\ 2\alpha_1 + 3\alpha_2 + 4\alpha_3 + 3\alpha_4 + 2\alpha_5 + 2\alpha_t + \alpha_{t'} & \text{in the case 4.} \end{cases}$$

Then we have $\gamma \in (\Phi^{\perp[y^{(i)}]})^+$, since $r_1 \notin [y^{(i)}]$ by the condition $m = N - l \geq 3$ mentioned above. We have $\gamma' = (w_x^s)^{-1} u'^{-1} \cdot \gamma \in (\Phi^{\perp[x]})^+$, $\gamma' \notin \Phi_J$ (since $u' w_x^s \cdot \Phi_J = \Phi_J$ and $\gamma \notin \Phi_J$) and $\langle \gamma', \alpha_1 \rangle \neq 0$ (since $\langle \gamma, \alpha_1 \rangle \neq 0$). The first and the third properties imply (since $\alpha_1 \in \Psi$ and Ψ is a union of irreducible components of $\Pi^{[x]}$) that γ' is a root in the root system of $(W(\Psi), S(\Psi))$, therefore γ' is a linear combination of elements of Ψ . This contradicts the second property $\gamma' \notin \Phi_J$. Hence these cases do not occur, as desired.

We show that the case 5 leads a contradiction. In this case, $y^{(i+1)}$ is obtained from $y^{(i)}$ by replacing r_4 and r_5 with r_5 and r_2 , respectively (note that $r_1, r_2 \notin [y^{(i)}]$ by the condition $m = N - l \geq 3$), and $J^{(i+1)} = \{r_2, r_3, r_4, r_5, t\}$. In other words, the first wide transformation ω maps the pair (r_4, r_5) to (r_5, r_2) and fixes the remaining points in $y^{(i)}$. Now by the same argument as above, the wide transformation in u next to ω (if it exists) is also of type 5 above (under a suitable renumbering of the elements) and it maps the pair (r_5, r_2) to either (r_4, r_5) or (r_2, r_4) . Iterating the argument, it follows that each wide transformation in u acts on the set $\{r_2, r_3, r_4, r_5\}$ as a rotation of three points r_2, r_4 and r_5 with fixed central point r_3 . This means that u cannot exchange the two points r_4 and r_5 as required by the property $u \in Y_{x, y}$ (recall that y was obtained from x by exchanging r_4 and r_5). Hence we have deduced a contradiction, as desired.

Finally, we show that the case 7 leads a contradiction. In this case, $y^{(i+1)}$ is obtained from $y^{(i)}$ by replacing r_4 with r_1 (note that $r_1, r_2 \notin [y^{(i)}]$ by the condition $m = N - l \geq 3$), and $J^{(i+1)} = J^{(i)} = J$. In other words, focusing on the three points r_1, r_3 , and r_4 , the wide transformation ω fixes r_3 that is adjacent to $t = t^{(i)}$, and maps r_4 to the remaining point r_1 . Iterating a similar argument, it follows that $J^{(h)} = J$ for every index h in the decomposition of u . Now we write $y^{(i)} = (r_3, r_4; y'^{(i)})$; note that $[y'^{(i)}] \cap J = \emptyset$ and $[y'^{(i)}]$ is not adjacent to J . Then we show that u contains at least one wide transformation $\omega' = \omega_{y'^{(h)}, J}^{t^{(h)}}$ of one of the following two forms:

- $y^{(h)}$ is of the form $y^{(h)} = (r_3, r_1; y'^{(h)})$, $t^{(h)}$ is adjacent to r_1 , and $\omega \cdot y^{(h)} = (r_4, r_1; y'^{(h)})$.
- $y^{(h)}$ is of the form $y^{(h)} = (r_1, r_4; y'^{(h)})$, $t^{(h)}$ is adjacent to r_1 , and $\omega \cdot y^{(h)} = (r_1, r_3; y'^{(h)})$.

Indeed, if such a wide transformation does not exist in u , it follows by iterating the above argument that the pair (r_3, r_4) is moved during the wide transformations in u only within the range $\{(r_3, r_1), (r_3, r_4), (r_1, r_4)\}$. Thus u cannot exchange the two points r_3 and r_4 as required by the property $u \in Y_{x,y}$ (recall that y was obtained from x by exchanging r_3 and r_4), a contradiction. Hence u contains at least one such wide transformation $\omega' = \omega_{y^{(h)}, J}^{t^{(h)}}$. Let h be the least index corresponding to such a wide transformation ω' . Put $t' = t^{(h)}$. Note that both $K_1 = \{t, r_3, r_2, r_1, r_4\}$ (the support of i -th transformation) and $K_2 = \{t', r_1, r_2, r_3, r_4\}$ (the support of h -th transformation) are of type D_5 , where the orderings of elements are according to the canonical labelling in Section 2.2.4.

Now for each transformation $\omega'' = \omega_{y^{(\nu)}, J}^{t^{(\nu)}}$ in u with $i < \nu < h$, if ω'' is a wide one then its action fixes every point in $[y^{(\nu)}] \setminus J$ (since ω'' is like in the case 7 above); while if ω'' is a narrow one then its action fixes every point in Π_J . This implies that the product w' (as elements of W) of all narrow transformations in u between $\omega = \omega_{y^{(i)}, J}^{t^{(i)}}$ and $\omega' = \omega_{y^{(h)}, J}^{t^{(h)}}$ (arranged in the same order) satisfies that $w' \in Y_{z, y^{(i)}}$ where $z = (r_3, r_4; y'^{(h)})$. Moreover, by the definition of supports of (narrow or wide) transformations, $[y'^{(i)}]$ is not adjacent to K_1 and $[y'^{(h)}]$ is not adjacent to K_2 . By the above properties, we have $\alpha_{t'} \in \Pi^{[z]}$ and $\gamma = w'' \cdot \alpha_{t'} \in \Pi^{[x]}$ where $w'' = (w' u' w_x^s)^{-1} \in Y_{x,z}$, while $w'' \cdot \alpha_1 = \beta_0$ and $w'' \cdot \beta_0 = \alpha_1$ (since both w' and u' fixes every point in Π_J , $w_x^s \cdot \beta_0 = \alpha_1$ and $w_x^s \cdot \alpha_1 = \beta_0$), therefore $\gamma \notin \Psi = \{\alpha_1, \beta_0\}$ and $\langle \gamma, \beta_0 \rangle = \langle \alpha_{t'}, \alpha_1 \rangle \neq 0$. This contradicts the definition of Ψ , as desired.

Hence we have deduced a contradiction in any of the above seven cases, concluding the proof for the first choice of $x_{(J)}$ in the case of type D_N .

From now, we consider the second choice of $x_{(J)}$, namely $x_{(J)}$ consists of the elements $r_{N-1}, r_{N-3}, \dots, r_{N-2k-1}$. Put $m = N-2k-1$. Now $\Psi = \Pi^{J, [x_{(J)}]}$ consists of α_j for $1 \leq j \leq m-2$, $\beta_i = \alpha_{m-2+2i} + \sum_{j=m-1+2i}^{N-2} 2\alpha_j + \alpha_{N-1} + \alpha_N$ for $0 \leq i \leq k$ (where β_0 is defined only when $m \geq 3$), and α_N . Note that $m \neq 2$ since $\Psi \cup \Pi_{[x_{(J)}]}$ spans V_J . First we consider the case $k \geq 1$, hence $r_{N-3} \in [x_{(J)}]$. Then in a similar way to Lemma 7.10, the facts $w \cdot \beta_k \in \Psi$, $w \cdot \alpha_N \in \Psi$ and

$$w \cdot \alpha_{N-2} = (w \cdot \beta_k - w \cdot \alpha_N - \alpha_{N-3} - \alpha_{N-1})/2 \in \Phi$$

imply that $\{w \cdot \beta_k, w \cdot \alpha_N\} = \{\beta_k, \alpha_N\}$ (namely, the last condition implies that the two coefficients in $w \cdot \beta_k$ and in $w \cdot \alpha_N$ of each of $\alpha_{m-2}, \alpha_m, \dots, \alpha_{N-5}, \alpha_N$ must have the same parity). Since $\{\beta_k\}$ and $\{\alpha_N\}$ form irreducible components of Ψ and $\langle w \rangle$ acts transitively on the set of irreducible components of Ψ , it follows that $\Psi = \{\beta_k, \alpha_N\}$, therefore we have $m = 1$ and $k = 1$, i.e., $N = 4$ and $x_{(J)}$ consists of r_1 and r_3 . By symmetry, this case is reduced to the case of the first choice of $x_{(J)}$ already discussed above.

On the other hand, we consider the case $k = 0$, hence $x_{(J)}$ consists of r_{N-1} only. In this case, $\Psi = \Pi^{J, [x_{(J)}]}$ consists of α_j for $1 \leq j \leq N-3$, $\beta = \alpha_{N-3} + 2\alpha_{N-2} + \alpha_{N-1} + \alpha_N$, and α_N . If $N \geq 5$, then Ψ is of type $D_{N-2} \times A_1$ (where $D_3 = A_3$), while all irreducible components of Ψ should be isomorphic since $\langle w \rangle$ acts transitively on the set of irreducible components of Ψ (see also Theorem 4.6(4)). This is a contradiction, therefore we have $N = 4$. Now by changing the representative of $x_{(J)}$, we may assume without loss of generality that $x_{(J)}$ consists of r_2 only. Then the action of w permutes the three irreducible components $\{\beta_i\}$, $1 \leq i \leq 3$ of Ψ cyclically, where $\beta_1 = \alpha_1 + \alpha_2 + \alpha_3$, $\beta_2 = \alpha_1 + \alpha_2 + \alpha_4$ and $\beta_3 = \alpha_2 + \alpha_3 + \alpha_4$. By symmetry, we may assume without loss of generality that $w \cdot \beta_1 = \beta_2$, $w \cdot \beta_2 = \beta_3$ and $w \cdot \beta_3 = \beta_1$. Then we have $w \cdot \alpha_1 = \alpha_4$, $w \cdot \alpha_2 = \alpha_2$, $w \cdot \alpha_3 = \alpha_1$ and $w \cdot \alpha_4 = \alpha_3$, therefore $w \cdot \Pi_J = \Pi_J$. Now we

consider a decomposition $w = \omega_{y^{(n)}, J^{(n)}}^{t^{(n)}} \cdots \omega_{y^{(1)}, J^{(1)}}^{t^{(1)}} \omega_{y^{(0)}, J^{(0)}}^{t^{(0)}}$ of w as in Section 7.3 with respect to $J^{(0)} = J$. Since w does not fix $\alpha_1 \in \Pi_J$, at least one transformation $\omega = \omega_{y^{(i)}, J^{(i)}}^{t^{(i)}}$ should be a wide one (otherwise w should fix Π_J pointwise). Let i be the least such index. Then we have $J^{(i)} = J$ and $[y^{(i)}] \cap J = \{r_2\}$. Now the support K of ω is irreducible and of finite type, $J \cup \{t^{(i)}\} \subseteq K$, $t^{(i)}$ is adjacent to $K \cap [y^{(i)}]$ in Γ , and $[y^{(i)}] \setminus J$ is not adjacent to J . These conditions imply that the unique possibility (up to symmetry) is that $t = t^{(i)}$ is adjacent to r_1 , some $t' \in [y^{(i)}] \setminus J$ is adjacent to t and $K = \{t', t, r_1, r_2, r_3, r_4\}$ is of type D_6 , where the ordering of elements is according to the canonical labelling in Section 2.2.4. However, now ω fixes every point in $\Pi_{K \setminus \{t\}}$, contradicting Lemma 7.6. Thus we have a contradiction in any case.

Hence the proof for the case of type D_N is concluded.

Before we continue the proof, we give a notation and some remarks as follows. First, for simplicity, we introduce an abbreviation

$$[c_1 c_2 \cdots c_N] = \sum_{j=1}^N c_j \alpha_j, \text{ where } N = |J|.$$

Secondly, note that we have no need to consider the case that $x_{(J)}$ consists of $|J| - 1$ or more elements, since then either Ψ is empty (a contradiction) or $|\Psi| = 1$ (forcing w to fix Ψ pointwise, as desired). In particular, we have no need to consider the case that J is of type $I_2(m)$.

Type E_6 First we consider the case that $x_{(J)}$ is of type A_1 . By considering a representative of $x_{(J)}$, we may assume without loss of generality that $x_{(J)} = (r_2)$. Then the set $\Psi = \Pi^{J, [x_{(J)}]} = \{\alpha_3, \alpha_1, [011210], \alpha_6, \alpha_5\}$ is of type A_5 , where the ordering of elements is canonical in Section 2.2.4 (similarly in the following argument, unless otherwise specified). Assume contrary that the action of w induces a nontrivial automorphism on Ψ , i.e., w exchanges α_3 and α_5 , exchanges α_1 and α_6 , and fixes $[011210]$. Then we have $w \cdot \alpha_4 = \alpha_4$, therefore $w \cdot \Pi_J = \Pi_J$. Now an argument similar to the case of type D_N can be applied. Namely, let $w = \omega_{y^{(n)}, J^{(n)}}^{t^{(n)}} \cdots \omega_{y^{(0)}, J^{(0)}}^{t^{(0)}}$ be a decomposition of w as in Section 7.3 with respect to $J^{(0)} = J$. Since w does not fix $\alpha_1 \in \Pi_J$, some of the transformations in w is a wide one; let $\omega = \omega_{y^{(i)}, J^{(i)}}^{t^{(i)}}$ be a wide transformation with least index i . Now by the shape of J , only the possibility (up to symmetry) is that the support K of ω satisfies that $K = \{r_1, r_2, \dots, r_6, t, t'\}$ is of type E_8 , where $t = t^{(i)}$ and $t' \in [y^{(i)}] \setminus J$. Then we have $\gamma = [011222] + 2\alpha_t + \alpha_{t'} \in (\Phi^{\perp[y^{(i)}]})^+$ and $\langle \gamma, \alpha_1 \rangle \neq 0$, therefore we have $\gamma' = u^{-1} \cdot \gamma \in (\Phi^{\perp[x]})^+ \setminus \Phi_J$ where $u = \omega_{y^{(i-1)}, J^{(i-1)}}^{t^{(i-1)}} \cdots \omega_{y^{(0)}, J^{(0)}}^{t^{(0)}}$, and $\langle \gamma', \alpha_1 \rangle \neq 0$ since $u \cdot \alpha_1 = \alpha_1$. Since $\Psi \subseteq \Phi_J$ and $\alpha_1 \in \Psi$, this contradicts the property (by definition) that Ψ is an irreducible component of $\Pi^{[x]}$. Thus the claim holds in this case.

Here we give a remark to shorten the following argument. Suppose that $x_{(J)}$ has an irreducible component of type A_1 , say $\{r\}$. Let $y \in S^*$ denote the remaining part of x ; $x = (r; y)$. Then we have $\Pi_{[y]} \subseteq \Pi^{\{r\}}$ and $\Phi_J^{\perp[x_{(J)}]} = (\Phi_J^{\perp\{r\}})^{\perp[y] \cap J}$, where

$$(\Phi_J^{\perp\{r\}})^{\perp[y] \cap J} = \{\gamma \in \Phi_J^{\perp\{r\}} \mid \langle \gamma, \alpha_s \rangle = 0 \text{ for every } s \in [y] \cap J\}.$$

Now since the root system $\Phi_J^{\perp\{r\}}$ for $W_J^{\perp\{r\}}$ is of type A_5 as above, the above argument for the case that J is of type A_N shows that the union of $\Pi_{[y] \cap J}$ and the simple system for $(\Phi_J^{\perp\{r\}})^{\perp[y] \cap J}$ cannot span the vector space spanned by $\Phi_J^{\perp\{r\}}$. This implies that the union of $\Pi_{[x_{(J)}]}$ and the simple system $\Pi^{J, [x_{(J)}]}$ for $\Phi_J^{\perp[x_{(J)}]}$ cannot span the vector space V_J , contradicting our assumption on Ψ . Thus we have no need to consider the case that $x_{(J)}$ has an irreducible component of type A_1 . The same argument works when for the union y of some (possibly a single) irreducible

components of $x_{(J)}$, we have known that $\Pi^{J,[y]}$ is of type A_n ; we have no need to consider the case of such an $x_{(J)}$. On the other hand, if for the union y of some (possibly a single) irreducible components of $x_{(J)}$ we have known that $\Pi^{J,[y]} \cup \Pi_{[y]}$ cannot span V_J , then it follows immediately that $\Pi^{J,[x_{(J)}]} \cup \Pi_{[x_{(J)}]}$ cannot span V_J as well, hence we have no need to consider such a case. Similar arguments will be used many times in the following argument as well.

By this remark and the assumption on the types of irreducible components of I , only the remaining case is that $x_{(J)}$ is of type D_4 , i.e., $x_{(J)}$ consists of r_2, r_3, r_4 , and r_5 . However, now we have $\Psi = \Pi^{J,[x_{(J)}]} = \emptyset$, a contradiction. Hence the proof for the case of type E_6 is concluded.

Type E_7 First we consider the case that $x_{(J)}$ is of type A_1 . We may assume without loss of generality that $x_{(J)} = (r_2)$. Then $\Psi = \Pi^{J,[x_{(J)}]} = \{\alpha_3, \alpha_1, [0112100], \alpha_6, \alpha_5, \alpha_7\}$ is of type D_6 , therefore if w induces a nontrivial automorphism on Ψ , then w exchanges α_5 and α_7 and fixes the other elements of Ψ . This implies that

$$w \cdot \alpha_4 = (w \cdot [0112100] - w \cdot [0110100])/2 = ([0112100] - [0110001])/2 \notin \Phi ,$$

a contradiction. Thus the claim holds in this case.

Secondly, we consider the case that $x_{(J)}$ is of type $A_1 \times A_1$. We may assume without loss of generality that $x_{(J)} = (r_2, r_7)$. Then $\Pi^{J,[x_{(J)}]}$ consists of only four elements $\alpha_1, \alpha_3, [0112100]$, and $[0112221]$, therefore $\Psi \cup \Pi_{[x_{(J)}]}$ cannot span V_J . Thus by the above remark, we have no need to consider the case that $x_{(J)}$ has two or more irreducible components of type A_1 .

Thus there are three remaining cases; $x_{(J)}$ is of type D_4, D_5 , and $D_4 \times A_1$. If $x_{(J)}$ is of type D_5 , then we may assume without loss of generality that $x_{(J)} = (r_1, r_2, r_3, r_4, r_5)$, and now we have $\Pi^{J,[x_{(J)}]} = \{\alpha_7\}$ and $\Pi^{J,[x_{(J)}]} \cup \Pi_{[x_{(J)}]}$ cannot span V_J . On the other hand, if $x_{(J)}$ is of type $D_4 \times A_1$, i.e., $x_{(J)} = (r_2, r_3, r_4, r_5, r_7)$, then we have $\Pi^{J,[x_{(J)}]} = \{[2234321]\}$ and $\Pi^{J,[x_{(J)}]} \cup \Pi_{[x_{(J)}]}$ cannot span V_J . Thus in fact the remaining case is only the case that $x_{(J)}$ is of type D_4 , i.e., $x_{(J)} = (r_2, r_3, r_4, r_5)$. Now $\Psi = \Pi^{J,[x_{(J)}]}$ consists of $\alpha_7, \beta = [0112221]$ and $\beta' = [2234321]$, and Ψ is of type $A_1 \times A_1 \times A_1$. By the facts $w \cdot \alpha_7 \in \Psi, w \cdot \beta \in \Psi$ and

$$w \cdot \alpha_6 = (w \cdot \beta - [0112200] - w \cdot \alpha_7)/2 \in \Phi ,$$

it follows that either w fixes α_7 and β , or w exchanges α_7 and β . In the former case, w fixes Ψ pointwise as desired. On the other hand, in the latter case, w fixes the remaining element β' and we have

$$w \cdot \alpha_1 = w \cdot (\beta' - \beta - [0122100])/2 = ([2234321] - [0122101])/2 \notin \Phi ,$$

a contradiction. Hence the proof for the case of type E_7 is concluded.

Type E_8 First we consider the case that $x_{(J)}$ is of type A_1 . We may assume without loss of generality that $x_{(J)} = (r_2)$. Then $\Psi = \Pi^{J,[x_{(J)}]} = \{\alpha_8, \alpha_5, \alpha_7, \alpha_6, [01121000], \alpha_1, \alpha_3\}$ is of type E_7 , therefore the automorphism on Ψ induced by w must be trivial. Hence w fixes Ψ pointwise, as desired.

We consider the case that $x_{(J)}$ is of type $A_1 \times A_1$. We may assume without loss of generality that $x_{(J)} = (r_2, r_8)$. Then $\Psi = \Pi^{J,[x_{(J)}]} = \{\alpha_5, \alpha_6, [01121000], \alpha_1, \alpha_3, [01122221]\}$ is of type D_6 . If w induces a nontrivial automorphism on Ψ , then w exchanges α_3 and $[01122221]$ and fixes the other elements of Ψ . Now we have

$$w \cdot \alpha_4 = w \cdot ([01121000] - [01101000])/2 = ([01121000] - [01001000] - [01122221])/2 \notin \Phi ,$$

a contradiction. Hence the claim holds in this case.

We consider the case that $x_{(J)}$ is of type $A_1 \times A_1 \times A_1$. We may assume without loss of generality that $x_{(J)} = (r_2, r_3, r_8)$. Then $\Psi = \Pi^{J, [x_{(J)}]}$ is the union of $\{[01122221]\}$ and $\{\alpha_5, \alpha_6, \beta, \beta'\}$, where $\beta = [01121000]$ and $\beta' = [22343221]$, and Ψ is of type $D_4 \times A_1$. This is a contradiction, since all irreducible components of Ψ should be isomorphic by the definition of Ψ .

We consider the case that $x_{(J)}$ is of type $A_1 \times A_1 \times A_1 \times A_1$. We may assume without loss of generality that $x_{(J)} = (r_2, r_3, r_5, r_8)$. Then $\Psi = \Pi^{J, [x_{(J)}]}$ consists of $[01121000]$, $[22343221]$, $[23465421]$, and $[01122221]$, and Ψ is of type $A_1 \times A_1 \times A_1 \times A_1$. Now since $w \cdot [01121000] \in \Psi$ and

$$w \cdot \alpha_4 = w \cdot ([01121000] - [01101000])/2 = (w \cdot [01121000] - [01101000])/2 \in \Phi ,$$

it follows that w fixes $[01121000]$. This contradicts the fact that $\langle w \rangle$ acts transitively on the set of irreducible components of Ψ .

We consider the case that $x_{(J)}$ is of type D_4 , i.e., $x_{(J)} = (r_2, r_3, r_4, r_5)$. Now $\Psi = \Pi^{J, [x_{(J)}]} = \{\alpha_7, \alpha_8, \beta, \beta'\}$ is of type D_4 , where $\beta = [01122210]$ and $\beta' = [22343210]$. By the shape of Ψ , any automorphism on Ψ fixes α_8 , therefore $w \cdot \alpha_8 = \alpha_8$ and $w \cdot (\Psi \setminus \{\alpha_8\}) = \Psi \setminus \{\alpha_8\}$. Now the restriction of the action of w to $V_{J \setminus \{r_8\}}$ is exactly the same as the one considered above for the case that J is of type E_7 and $x_{(J)}$ is of type D_4 . Thus by the previous argument, w fixes $\Psi \setminus \{\alpha_8\}$, hence Ψ , pointwise. Hence the claim holds in this case.

We consider the case that $x_{(J)}$ is of type D_5 . We may assume without loss of generality that $x_{(J)} = (r_1, r_2, r_3, r_4, r_5)$. Now $\Psi = \Pi^{J, [x_{(J)}]} = \{\alpha_8, \alpha_7, \beta\}$ is of type A_3 , where $\beta = [23465421]$. If w induces a nontrivial automorphism on Ψ , then w fixes α_7 and exchanges α_8 and β . Now

$$w \cdot \alpha_6 = w \cdot (\beta - [23465021])/4 = (\alpha_8 - [23465020] - \beta)/4 \notin \Phi ,$$

a contradiction. Hence w fixes Ψ pointwise, as desired. Since Ψ is now of type A_3 , by the above remark we have no need to consider the case that $x_{(J)}$ is of type $D_5 \times A_1$.

We consider the case that $x_{(J)}$ is of type $D_4 \times A_1$. We may assume without loss of generality that $x_{(J)} = (r_2, r_3, r_4, r_5, r_7)$. Now $\Psi = \Pi^{J, [x_{(J)}]}$ consists of three elements $\beta_1 = [01122210]$, $\beta_2 = [22343210]$ and $\beta_3 = [23465432]$, and Ψ is of type $A_1 \times A_1 \times A_1$. Now since $w \cdot \beta_1 \in \Psi$ and

$$w \cdot \alpha_6 = w \cdot (\beta_1 - [01122010])/2 = (w \cdot \beta_1 - [01122010])/2 \in \Phi ,$$

it follows that w fixes β_1 . This contradicts the fact that $\langle w \rangle$ acts transitively on the set of irreducible components of Ψ .

We consider the case that $x_{(J)}$ is of type D_6 , i.e., $x_{(J)} = (r_2, r_3, r_4, r_5, r_6, r_7)$. Now $\Psi = \Pi^{J, [x_{(J)}]}$ consists of $\beta = [22343210]$ and $\beta' = [23465432]$, and Ψ is of type $A_1 \times A_1$. Since $\langle w \rangle$ acts transitively on the set of irreducible components of Ψ , w must exchange β and β' . However, now we have

$$w \cdot \alpha_1 = w \cdot (\beta - [02343210])/2 = (\beta' - [02343210])/2 \notin \Phi ,$$

a contradiction.

Now only the remaining case is that $x_{(J)}$ is of type E_6 , i.e., $x_{(J)} = (r_1, r_2, r_3, r_4, r_5, r_6)$. Now $\Psi = \Pi^{J, [x_{(J)}]} = \{\alpha_8, [23465431]\}$ is of type A_2 . If w induces a nontrivial automorphism on Ψ , then w exchanges α_8 and $[23465431]$. However, now we have

$$w \cdot \alpha_7 = w \cdot ([23465431] - [23465401])/3 = (\alpha_8 - [23465400] - [23465431])/3 \notin \Phi ,$$

a contradiction. Thus w fixes the set Ψ pointwise, as desired.

Hence the proof for the case of type E_8 is concluded.

Type F_4 First we consider the case that $x_{(J)}$ is of type A_1 . By symmetry, we may assume without loss of generality that $x_{(J)} = (r_1)$. Then $\Psi = \Pi^{J, [x_{(J)}]} = \{\alpha_3, \alpha_4, [12\sqrt{2}0]\}$ is of type B_3 . Thus the automorphism on Ψ induced by w must be trivial, therefore w fixes Ψ pointwise, as desired. Hence the claim holds in this case.

Secondly, we consider the case that $x_{(J)}$ is of type $A_1 \times A_1$. We may assume without loss of generality that $x_{(J)} = (r_1, r_4)$. Then $\Psi = \Pi^{J, [x_{(J)}]}$ consists of $\beta = [12(2\sqrt{2})\sqrt{2}]$ and $\beta' = [\sqrt{2}(2\sqrt{2})21]$, and Ψ is of type $A_1 \times A_1$. Since $\langle w \rangle$ acts transitively on the set of irreducible components of Ψ , w must exchange β and β' . However, now we have

$$w \cdot \alpha_3 = w \cdot (\sqrt{2}\beta - \beta' - \alpha_4)/2 = (\sqrt{2}\beta' - \beta - \alpha_4)/2 = [120(-1)]/2 \notin \Phi ,$$

a contradiction.

Finally, we consider the case that $x_{(J)}$ is of type B_2 , i.e., $x_{(J)} = (r_2, r_3)$. Now $\Psi = \Pi^{J, [x_{(J)}]} = \{\beta, \beta'\}$ is of type B_2 , where $\beta = [01\sqrt{2}\sqrt{2}]$ and $\beta' = [\sqrt{2}\sqrt{2}10]$. If w induces a nontrivial automorphism on Ψ , then w exchanges β and β' . However, now we have

$$w \cdot \alpha_4 = w \cdot (\beta - [01\sqrt{2}0])/\sqrt{2} = (\beta' - [01\sqrt{2}0])/\sqrt{2} \notin \Phi ,$$

a contradiction. Thus w fixes Ψ pointwise, as desired.

Hence the proof for the case of type F_4 is concluded.

In the following proof for the case of type H_3 and H_4 , we put $c = (1 + \sqrt{5})/2$, therefore we have $c^2 = c + 1$, $r_1 \cdot \alpha_2 = c\alpha_1 + \alpha_2$ and $r_2 \cdot \alpha_1 = \alpha_1 + c\alpha_2$.

Type H_3 In this case, it suffices to consider the case that $x_{(J)}$ is of type A_1 . We may assume without loss of generality that $x_{(J)} = (r_1)$. Then the set $\Psi = \Pi^{J, [x_{(J)}]}$ consists of α_3 and $\beta = [(c+1)(2c)c]$, and Ψ is of type $A_1 \times A_1$. Since $\langle w \rangle$ acts transitively on the set of irreducible components of Ψ , w must exchange α_3 and β . However, now we have

$$w \cdot \alpha_2 = w \cdot (\beta - [(c+1)0c])/(2c) = (\alpha_3 - (c+1)\alpha_1 - c\beta)/(2c) \notin \Phi ,$$

a contradiction. Hence the proof for the case of type H_3 is concluded.

Type H_4 First we consider the case that $x_{(J)}$ is of type A_1 . We may assume without loss of generality that $x_{(J)} = (r_1)$. Then $\Psi = \Pi^{J, [x_{(J)}]} = \{\beta, \alpha_4, \alpha_3\}$ is of type H_3 , where $\beta = [(c+1)(2c)c0]$ (see the second last paragraph for the definition of c). Now the automorphism on Ψ induced by w must be trivial, therefore w fixes Ψ pointwise, as desired. Hence the claim holds in this case.

Secondly, we consider the case that $x_{(J)}$ is of type $A_1 \times A_1$. We may assume without loss of generality that $x_{(J)} = (r_1, r_3)$. Then $\Psi = \Pi^{J, [x_{(J)}]}$ consists of $\beta = [(c+1)(2c)c0]$ and $\beta' = [(3c+2)(4c+2)(3c+1)(2c)]$, and Ψ is of type $A_1 \times A_1$. Since $\langle w \rangle$ acts transitively on the set of irreducible components of Ψ , w must exchange β and β' . However, now we have

$$w \cdot \alpha_2 = w \cdot (\beta - [(c+1)0c0])/(2c) = (\beta' - [(c+1)0c0])/(2c) \notin \Phi ,$$

a contradiction.

Finally, we consider the case that $x_{(J)}$ is of type $I_2(5)$, i.e., $x_{(J)} = (r_1, r_2)$. Then $\Psi = \Pi^{J, [x_{(J)}]} = \{\alpha_4, \beta\}$ is of type $I_2(5)$, where $\beta = [(2c+1)(2c+2)(c+2)1]$. If w induces a nontrivial automorphism on Ψ , then w must exchange α_4 and β . However, now we have

$$w \cdot \alpha_3 = w \cdot (\beta - [(2c+1)(2c+2)01])/(c+2) = (\alpha_4 - [(2c+1)(2c+2)00] - \beta)/(c+2) \notin \Phi ,$$

a contradiction. Thus w fixes the set Ψ pointwise, as desired.

Hence the proof for the case of type H_5 is concluded.

Summarizing, the claim of Theorem 7.1 holds in the case $\Psi \not\subseteq \Phi_{I^\perp}$.

7.5 The second case $\Psi \subseteq \Phi_{I^\perp}$

In this subsection, we consider the remaining case that $\Psi \subseteq \Phi_{I^\perp}$ (see (4.14) for the definition of $I^\perp$). Note that $\Psi = \Pi_{J \setminus I}$ in this case, since $\Pi_{I^\perp} \subseteq \Pi^I$. Let L be a subset of S such that Π_L is an irreducible component of Ψ . In particular, L itself is irreducible and of finite type. The aim of the following argument is to show that w fixes Π_L pointwise. Indeed, if this is satisfied, then we have $\Psi = \Pi_L$ since $\langle w \rangle$ acts transitively on the set of irreducible components of Ψ (by the definition of Ψ) and w fixes Ψ pointwise, as desired. Put $x = x_I$. Note that $w \cdot \Pi_L \subseteq \Pi$ since now $\Psi \subseteq \Pi$ as mentioned above.

7.5.1 The possibilities of transformations

In this subsection, we determine the possibilities of narrow and wide transformations $\omega = \omega_{y^{(h)}, J^{(h)}}^{t^{(h)}}$ in a decomposition of w as in Section 7.3 with respect to $J^{(0)} = L$. Let K denote the support of ω , which is irreducible and of finite type. Here we use the canonical labelling $K = \{r_1, r_2, \dots, r_n\}$ on K given in Section 2.2.4. Put $y = y^{(h)}$, $L' = J^{(h)}$ and $t = t^{(h)}$. Then $t \in K$, and t is adjacent to an element of $[y] \cap K$.

First we consider the case of narrow transformations, therefore $L' \cap K = \emptyset$ and $K \setminus \{t\} \subseteq [y]$. Now recall that ω cannot fix the set $\Pi_{K \setminus \{t\}}$ pointwise (see Lemma 7.6). By the assumption on the type of irreducible components of I (i.e., it is not of type A_N , $N \geq 2$), the possibilities of K and t are determined (up to symmetry) as follows:

1. K is of type $I_2(m)$ with m odd, where $I_2(3) = A_2$, and $t = r_2$; now $\omega : r_1 \mapsto r_2$.
2. K is of type A_3 and $t = r_2$; now $\omega : (r_1, r_3) \mapsto (r_3, r_1)$.
3. K is of type D_n ($n \geq 5$) and $t = r_1$; now ω exchanges r_{n-1} and r_n and fixes every r_j , $2 \leq j \leq n-2$.
4. K is of type E_6 and $t = r_1$; now $\omega : (r_2, r_3, r_4, r_5, r_6) \mapsto (r_5, r_2, r_4, r_3, r_1)$.
5. K is of type E_7 and $t = r_6$; now $\omega : (r_1, r_2, r_3, r_4, r_5, r_7) \mapsto (r_1, r_5, r_3, r_4, r_2, r_7)$.
6. K is of type E_7 and $t = r_7$; now $\omega : (r_1, r_2, r_3, r_4, r_5, r_6) \mapsto (r_6, r_2, r_5, r_4, r_3, r_1)$.
7. K is of type E_8 and $t = r_1$; now $\omega : (r_2, r_3, r_4, r_5, r_6, r_7, r_8) \mapsto (r_3, r_2, r_4, r_5, r_6, r_7, r_8)$.
8. K is of type E_8 and $t = r_7$; now $\omega : (r_1, r_2, r_3, r_4, r_5, r_6, r_8) \mapsto (r_6, r_2, r_5, r_4, r_3, r_1, r_8)$.
9. K is of type H_3 and $t = r_3$; now $\omega : (r_1, r_2) \mapsto (r_2, r_1)$.
10. K is of type H_4 and $t = r_3$; now $\omega : (r_1, r_2, r_4) \mapsto (r_2, r_1, r_4)$.

An important property derived from the enumeration is the following:

Lemma 7.11. *For any narrow transformation as above, the union of the irreducible components of $K \setminus \{t\}$ of type A_1 is left invariant under the action of ω unless ω is in the case 1. Moreover, if it is not in the case 1 and $K \setminus \{t\}$ has an irreducible component of type A_1 , then the irreducible component of $K \setminus \{t\}$ of type other than A_1 (if it exists) is left invariant under the action of ω .*

Secondly, we consider the case of wide transformations, therefore we have $L' \subseteq K$ and $K \setminus (L' \cup \{t\}) \subseteq [y]$. Note that t is adjacent to $[y] \cap K$. Now recall that ω cannot fix the set $\Pi_{K \setminus \{t\}}$ pointwise (see Lemma 7.6). Moreover, it must hold that there does not exist a $\gamma \in (\Phi_K^{\perp[y] \cap K})^+$ such that $\gamma \notin \Phi_{K \setminus \{t\}}$. Indeed, if such a γ exists, then we have $\omega \cdot \gamma \in \Phi^-$ and

$\Phi^{\perp[y]}[\omega] \neq \emptyset$, while ω is a right divisor of $wu^{-1} \in Y_{x,y}$ where $u = \omega_{y^{(h-1)}, J^{(h-1)}}^{t^{(h-1)}} \cdots \omega_{y^{(0)}, J^{(0)}}^{t^{(0)}} \in Y_{y,x}$; this contradicts Lemma 2.2 and the definition of Y . On the other hand, L' is not adjacent to $[y] \cap K$, since we have $u \cdot \Pi_L = \Pi_{L'}$, $u \cdot \Pi_{[x]} = \Pi_{[y]}$, and $[x]$ and L are not adjacent in Γ . By these conditions and the assumption on the type of irreducible components of I , the possibilities are determined (up to symmetry) as follows:

1. K is of type A_n ($n \geq 3$), $t = r_2$ and $L' = \{r_3, \dots, r_n\}$ (type A_{n-2}). Now $\omega : r_1 \mapsto r_n$ and $\omega : (r_3, r_4, \dots, r_n) \mapsto (r_1, r_2, \dots, r_{n-2})$.
2. K is of type E_7 , $t = r_6$ and $L' = \{r_7\}$ (type A_1). Now $\omega : (r_1, r_2, r_3, r_4, r_5) \mapsto (r_1, r_5, r_3, r_4, r_2)$ and $\omega : r_7 \mapsto r_7$.

This implies the following property:

Lemma 7.12. *For any wide transformation as above, $[y] \cap K$ is irreducible and L' is of type A_N , $N \geq 1$. Moreover, if $[y] \cap K$ is not of type A_1 , then ω fixes $\Pi_{L'}$ pointwise and leaves the set $\Pi_{[y] \cap K}$ invariant.*

By virtue of this lemma, we can prove the following:

Lemma 7.13. *If L is not of type A_N , then w fixes the set Π_L pointwise, hence Theorem 7.1 holds in this case.*

Proof. Let $w = \omega_{y^{(h)}, J^{(h)}}^{t^{(h)}} \cdots \omega_{y^{(1)}, J^{(1)}}^{t^{(1)}} \omega_{y^{(0)}, J^{(0)}}^{t^{(0)}}$ be a decomposition of $w \in Y_{x,x}$ as in Section 7.3 with respect to $J^{(0)} = L$. Put $\omega_i = \omega_{y^{(i)}, J^{(i)}}^{t^{(i)}}$ for each i . Then we prove inductively that every transformation ω_i is a narrow one and it fixes the set Π_L pointwise. If this has been proven for $\omega_0, \omega_1, \dots, \omega_{i-1}$, then we have $J^{(i)} = L$, therefore Lemma 7.12 implies that ω_i cannot be a wide one. Thus ω_i is a narrow one, therefore ω_i fixes $\Pi_{J^{(i)}} = \Pi_L$ pointwise. This implies that the claim for the induction holds, therefore $w = \omega_h \cdots \omega_0$ fixes Π_L pointwise, as desired. $\square$

From now, we suppose that L is of type A_N . We divide the following argument into the two cases; the case $N \geq 2$ discussed in Section 7.5.2; and the case $N = 1$ discussed in Section 7.5.3.

7.5.2 Proof for the case $N \geq 2$

In this subsection, we suppose that L is of type A_N , $N \geq 2$, and prove that w fixes Π_L pointwise. Let $w = \omega_n \cdots \omega_1 \omega_0$, where $\omega_i = \omega_{y^{(i)}, J^{(i)}}^{t^{(i)}}$ for each i , be a decomposition of $w \in Y_{x,x}$ as in Section 7.3 with respect to $J^{(0)} = L$. First note that if the claim holds for $w_i = \omega_{i-1} \cdots \omega_1 \omega_0 \omega_n \cdots \omega_i \in Y_{y^{(i)}, y^{(i)}}$ and $\Pi_{J^{(i)}} = \omega_{i-1} \cdots \omega_1 \omega_0 \cdot \Pi_L$ instead of w and Π_L , then the original claim for w and Π_L also holds (i.e., w fixes Π_L pointwise). Thus if $\ell(w_i) < \ell(w)$, then the length of the element w can be decreased by considering w_i instead of w . Since this process terminates in a finite number of steps, we may assume without loss of generality that $\ell(w_i) = \ell(w)$ for every i (note that $\ell(w) = \sum_{j=0}^n \ell(\omega_j)$ by the definition of the decomposition, therefore $\ell(w_i) \leq \ell(w)$). This implies that $\ell(w_i) = \sum_{j=0}^n \ell(\omega_j)$, therefore the expression $w_i = \omega_{i-1} \cdots \omega_1 \omega_0 \omega_n \cdots \omega_i$ is also a decomposition of w_i as in Section 7.3. By a similar reason, $w_i^{-1} = \omega_i^{-1} \cdots \omega_n^{-1} \omega_0^{-1} \omega_1^{-1} \cdots \omega_{i-1}^{-1}$ is also a decomposition of w_i^{-1} as in Section 7.3 (note that ω_j^{-1} is also a wide or narrow transformation by the definition of ω_j). These properties will be used in the following argument many times.

The condition for the type of L implies that if ω_i is a wide transformation, then ω_i is of type 1 among the two types in Section 7.5.1. Let K_i denote the support of ω_i . From now, we assume contrary that w does not fix Π_L pointwise. Then at least one ω_i should be a wide

transformation: By considering $\omega_{i-1} \cdots \omega_1 \omega_0 \omega_n \cdots \omega_i$ instead of w if necessary, we may assume without loss of generality that ω_0 is a wide transformation. From now, for $i \geq 0$, we determine inductively subsets M_i of S and integers η_i , μ_i , $\lambda_{i,h}$ and $\rho_{i,h}$ ($0 \leq h \leq i$) such that $M_{i-1} \subseteq M_i$, $\mu_i \geq 0$, $\lambda_{i,h} \geq 0$, $\rho_{i,h} \geq 0$, and the following conditions are satisfied (see Figure 5 for an illustration):

1. $M_i = \{r_{\eta_i+j}\}_{j=0}^{\mu_i}$ and $m_{r_j, r_{j+1}} = 3$ for every $\eta_i \leq j \leq \eta_i + \mu_i - 1$.
2. For each $0 \leq h \leq i$, we have $[y^{(h)}] \cap M_i = \{r_{\eta_i+2j}\}_{j=0}^{\lambda_{i,h}-1} \cup \{r_{\eta_i+2\lambda_{i,h}+N+1+2j}\}_{j=0}^{\rho_{i,h}-1}$, $J^{(h)} = \{r_{\eta_i+2\lambda_{i,h}+j}\}_{j=0}^{N-1}$, $\mu_i = 2\lambda_{i,h} + N - 1 + 2\rho_{i,h}$, and $[y^{(h)}] \cap M_i$ and $[y^{(h)}] \setminus M_i$ are not adjacent to each other in Γ .
3. We have $|\lambda_{i,1} - \lambda_{i,0}| = 1$ (when $i \geq 1$), and for each $2 \leq h \leq i$, we have $|\lambda_{i,h} - \lambda_{i,h-1}| \leq 1$ (hence $|\rho_{i,h} - \rho_{i,h-1}| \leq 1$). Moreover, for each $1 \leq h \leq i$,
 - if $\lambda_{i,h} - \lambda_{i,h-1} = 1$, then ω_{h-1} is a wide transformation with support $K_{h-1} = \{r_{\eta_i+2\lambda_{i,h-1}+j}\}_{j=0}^{N+1} \supseteq J^{(h-1)}$ and $t^{(h-1)} = r_{\eta_i+2\lambda_{i,h-1}+N}$;
 - if $\lambda_{i,h} - \lambda_{i,h-1} = -1$, then ω_{h-1} is a wide transformation with support $K_{h-1} = \{r_{\eta_i+2\lambda_{i,h-1}+j}\}_{j=-2}^{N-1} \supseteq J^{(h-1)}$ and $t^{(h-1)} = r_{\eta_i+2\lambda_{i,h-1}-1}$;
 - if $\lambda_{i,h} - \lambda_{i,h-1} = 0$, then ω_{h-1} is a narrow transformation.
4. For any $j \in \{0, 1, \dots, (\mu_i - N + 1)/2\}$, there exists some $0 \leq h \leq i$ such that $\lambda_{i,h} = j$.




Figure 5: Example of the data M_i (here black circles signify elements of $y^{(h)}$ or $y^{(h+1)}$, respectively, and r_{η_i+j} is abbreviated to $+j$)

For the case $i = 0$, we define $M_0 = L = \{r_1, r_2, \dots, r_N\}$ which is of type A_N (where the labelling is canonical in Section 2.2.4), $\eta_0 = 1$, $\mu_i = N - 1$, and $\lambda_{i,0} = \rho_{i,0} = 0$. Note that these data indeed satisfy the requirements.

Now suppose that $i \leq n$ and we have determined such a subset M_i and associated parameters η_i , μ_i , $\lambda_{i,h}$ and $\rho_{i,h}$. We investigate the possibilities of the next transformation ω_i . From now, put $\alpha_j = \alpha_{r_j}$ for simplicity. First, we prove the following:

Lemma 7.14. *In the setting, the following situation does not occur: ω_i is a narrow transformation of type 1 among the ten types in Section 7.5.1 whose support K_i contains an element of $[y^{(i)}] \cap M_i$.*

Proof. Assume contrary that ω_i is a narrow transformation of type 1 whose support K_i contains an element of $[y^{(i)}] \cap M_i$. By symmetry, we may assume without loss of generality that K_i contains an element $r_{\eta_i+2j_0}$ of $[y^{(i)}] \cap M_i$ with $0 \leq j_0 \leq \lambda_{i,i} - 1$. By the definition of narrow transformations of type 1, we have $K_i = \{r_{\eta_i+2j_0}, s\}$ for some $s \in S \setminus ([y^{(i)}] \cup J^{(i)})$ and $m_{r_{\eta_i+2j_0}, s}$

is odd. Now s is not adjacent to $([y^{(i)}] \cup J^{(i)}) \setminus \{r_{\eta_i+2j_0}\}$, hence $s \notin M_i$. By the condition 4 for M_i , we have $\lambda_{i,h} = j_0$ for some $0 \leq h < i$, therefore $J^{(h)} = \{r_{\eta_i+2j_0+j'}\}_{j'=0}^{N-1}$. Choose the largest such index h . Then since $\lambda_{i,h} < \lambda_{i,i}$, the maximality of h and the condition 3 for M_i imply that $\lambda_{i,h+1} = \lambda_{i,h} + 1$, ω_h is a wide transformation with support $K_h = \{r_{\eta_i+2j_0+j'}\}_{j'=0}^{N+1}$ and $\lambda_{i,h'} \geq j_0 + 1$ for every $h+1 < h' < i$. From now, we show that there exists a $\gamma \in \Phi^{\perp[y^{(h)}]}$ such that $\gamma \notin \Phi_{J^{(h)}}$ and γ is not orthogonal to $\alpha_{\eta_i+2j_0} \in \Pi_{J^{(h)}}$; if this is satisfied, then $(\omega_{h-1} \cdots \omega_1 \omega_0)^{-1} \cdot \gamma$ belongs to $\Phi^{\perp[x]} \setminus \Phi_L$ (note that $y^{(0)} = x$ and $J^{(0)} = L$) and is not orthogonal to $\alpha_{\eta_i+2\lambda_{i,0}} = (\omega_{h-1} \cdots \omega_1 \omega_0)^{-1} \cdot \alpha_{\eta_i+2j_0} \in \Pi_L$ (see the condition 3 for M_i), contradicting the fact that Π_L is an irreducible component of $\Pi^{[x]}$.

By the conditions for M_i and the above property derived from the maximality of h , each $\omega_{h'}$ ($h+1 \leq h' < i$) is one of the following three types:

- A wide transformation with support $K_{h'} \subseteq \{r_{\eta_i+j'} \mid j' \geq 2j_0 + 2\}$.
- A narrow transformation with support $K_{h'}$ containing no elements of $[y^{(h')}] \cap M_i$.
- A narrow transformation of not of type 1 in Section 7.5.1 with support $K_{h'}$ containing some element of $[y^{(h')}] \cap M_i$.

Let $h+1 < h'_1 < h'_2 < \cdots < h'_k < i$ be the indices h' such that $\omega_{h'}$ is of the third type. Then by Lemma 7.11 and the shape of $[y^{(h')}] \cap M_i$, $\omega_{h'_\nu}$ leaves the sets $[y^{(h'_\nu)}]$ and $\Pi_{J^{(h'_\nu)}}$ invariant for each $1 \leq \nu \leq k$. Now by applying Lemma 7.7 to $u = \omega_{i-1} \cdots \omega_{h'_k} \cdots \omega_{h'_1} \cdots \omega_{h+1}$, it follows that there is a $z \in \mathcal{V}(C)$ such that $[z] = [y^{(i)}]$ and $u \in Y_{z,y^{(h+1)}}$. Moreover, let $\lambda \in \Lambda$ be the index such that $(y^{(h+1)})_\lambda = r_{\eta_i+2j_0}$. Then the above construction implies that the support of every ω_j appearing in u does not contain $(y^{(h+1)})_\lambda$, and $u \cdot \alpha_{\eta_i+2j_0+N+1} = \alpha_{\eta_i+2\lambda_{i,i}+N-1}$. By Lemma 7.8, we have $u \in \widehat{Y}_{z',y'}$, where y' and z' are obtained from $y^{(h+1)}$ and z by replacing $(y^{(h+1)})_\lambda = z_\lambda$ with $r_{\eta_i+2j_0+N+1} \in J^{(h+1)}$ and with $r_{\eta_i+2\lambda_{i,i}+N-1} \in J^{(i)}$, respectively. Note that $y' = y^{(h)}$ by the property of ω_h , therefore $z' \in \mathcal{V}(C)$ and $u \in Y_{z',y^{(h)}}$. Now recall that $s \in S \setminus M_i$ is adjacent to $r_{\eta_i+2j_0}$ and is not adjacent to $([y^{(i)}] \cup J^{(i)}) \setminus \{r_{\eta_i+2j_0}\}$. By the choice of z' , it follows that s is not adjacent to $[z']$, therefore $\alpha_s \in \Pi^{[z']}$. Moreover, by the construction of u , the support of each ω_j appearing in u does not contain s or $r_{\eta_i+2j_0}$. This implies that the coefficients of α_s and $\alpha_{\eta_i+2j_0}$ in $\gamma = u^{-1} \cdot \alpha_s \in \Pi^{[y^{(h)}]}$ are equal to those of α_s and $\alpha_{\eta_i+2j_0}$ in α_s , namely 1 and 0, respectively. Thus γ satisfies the above requirement, hence the proof of Lemma 7.14 is concluded. $\square$

By Lemma 7.14, if ω_i is a narrow transformation, then either its support contains no elements of $[y^{(i)}] \cap M_i$ or it is not of type 1. Now Lemma 7.11 implies that, in any case, the choices $M_{i+1} = M_i$, $\eta_{i+1} = \eta_i$, $\mu_{i+1} = \mu_i$, $\lambda_{i+1,h} = \lambda_{i,h}$, $\rho_{i+1,h} = \rho_{i,h}$ ($0 \leq h \leq i$), $\lambda_{i+1,i+1} = \lambda_{i,i}$ and $\rho_{i+1,i+1} = \rho_{i,i}$ satisfy the conditions for M_{i+1} .

From now, we consider the case that ω_i is a wide transformation. Let K_i denote the support of ω_i . Let t and t' be the unique elements of $K_i \setminus ([y^{(i)}] \cup J^{(i)})$ and of $K_i \cap [y^{(i)}]$, respectively. If $t \in M_i$, then the shape of M_i implies that either $t = r_{\eta_i+2\lambda_{i,i}-1}$ and $t' = r_{\eta_i+2\lambda_{i,i}-2}$, or $t = r_{\eta_i+2\lambda_{i,i}+N}$ and $t' = r_{\eta_i+2\lambda_{i,i}+N+1}$. In the former case, the action of ω_i maps t' and $J^{(i)}$ to $r_{\eta_i+2\lambda_{i,i}+N-1}$ and $\{r_{\eta_i+2\lambda_{i,i}-2+j'}\}_{j'=0}^{N-1}$, respectively, therefore $M_{i+1} = M_i$, $\eta_{i+1} = \eta_i$, $\mu_{i+1} = \mu_i$, $\lambda_{i+1,h} = \lambda_{i,h}$, $\rho_{i+1,h} = \rho_{i,h}$ ($0 \leq h \leq i$), $\lambda_{i+1,i+1} = \lambda_{i,i} - 1$ and $\rho_{i+1,i+1} = \rho_{i,i} + 1$ satisfy the conditions for M_{i+1} . The latter case is similar. Thus we may assume that $t \notin M_i$. Note that t is adjacent to either $r_{\eta_i+2\lambda_{i,i}} \in J^{(i)}$ or $r_{\eta_i+2\lambda_{i,i}+N-1} \in J^{(i)}$. By symmetry, we may assume without loss of generality that t is adjacent to $r_{\eta_i+2\lambda_{i,i}}$. First, we have the following result:

Lemma 7.15. *In the above situation, the case $\lambda_{i,i} \geq 1$ and $t' = r_{\eta_i+2\lambda_{i,i}-2}$ does not occur.*

Proof. Assume contrary that $\lambda_{i,i} \geq 1$ and $t' = r_{\eta_i+2\lambda_{i,i}-2}$. Let h be the largest index $0 \leq h < i$ such that $\lambda_{i,h} = \lambda_{i,i} - 1$. Let $\lambda \in \Lambda$ be the index such that $(y^{(h+1)})_\lambda = t'$. Then by the condition 3 for M_i , we have $\lambda_{i,h+1} = \lambda_{i,i}$ and $\lambda_{i,h'} \geq \lambda_{i,i}$ for every $h+1 < h' < i$. Now by an argument similar to Lemma 7.14, each $\omega_{h'}$ ($h+1 \leq h' < i$) is one of the following three types:

- A wide transformation with support $K_{h'} \subseteq \{r_{\eta_i+j'} \mid j' \geq 2\lambda_{i,i} + 2\}$.
- A narrow transformation with support $K_{h'}$ containing no elements of $[y^{(h')}] \cap M_i$.
- A narrow transformation of not of type 1 in Section 7.5.1 with support $K_{h'}$ containing some element of $[y^{(h')}] \cap M_i$.

Then by an argument similar to Lemma 7.14 (using Lemmas 7.7 and 7.8), there exist a $z \in S^{(\Lambda)}$ and an element u such that $[z] = [y^{(i)}]$, $u \in Y_{z,y^{(h+1)}}$, $u \cdot \alpha_{\eta_i+2\lambda_{i,i}+N-1} = \alpha_{\eta_i+2\lambda_{i,i}+N-1}$, u is a product of wide or narrow transformations with support not containing $t' = (y^{(h+1)})_\lambda$ (hence not containing t), and $u \in \widehat{Y}_{z',y'}$, where y' and z' are obtained from $y^{(h+1)}$ and z , respectively, by replacing $(y^{(h+1)})_\lambda = z_\lambda = t'$ with $r_{\eta_i+2\lambda_{i,i}+N-1} \in J^{(h+1)} = J^{(i)}$. By the choice of h and the condition 3 for M_i , we have $y' = y^{(h)}$, therefore $u \in Y_{z',y^{(h)}}$. Now by the property of the wide transformation ω_i mentioned above, t is not adjacent to $[y^{(i)}] \setminus \{t'\}$, and t is not adjacent to $r_{\eta_i+2\lambda_{i,i}+N-1}$ since the support K_i of ω_i is of type A_{N+2} and $N \geq 2$. This implies that t is not adjacent to $[z']$ and $\alpha_t \in \Pi^{[z']}$, therefore $\gamma \in u^{-1} \cdot \alpha_t \in \Pi^{[y^{(h)}]}$. Moreover, by the above property of u , the coefficients of α_t and $\alpha_{t'}$ in γ are equal to those of α_t and $\alpha_{t'}$, respectively, in α_t ; namely 1 and 0. This implies that $\gamma \in \Pi^{[y^{(h)}]} \setminus \Phi_{J^{(h)}}$ is not orthogonal to $\alpha_{t'} \in \Pi_{J^{(h)}}$. In the same way as Lemma 7.14, this leads a contradiction that $(\omega_{h-1} \cdots \omega_1 \omega_0)^{-1} \cdot \gamma \in \Pi^{[x]} \setminus \Pi_L$ is not orthogonal to an element of Π_L . Hence the proof of Lemma 7.15 is concluded. $\square$

Secondly, we have the following result:

Lemma 7.16. *In the above situation, the case $\lambda_{i,i} \geq 1$ and $t' \notin \{r_{\eta_i+2\lambda_{i,i}+N+1+2j}\}_{j=0}^{\rho_{i,i}-1}$ does not occur.*

Proof. Assume contrary that $\lambda_{i,i} \geq 1$ and $t' \notin \{r_{\eta_i+2\lambda_{i,i}+N+1+2j}\}_{j=0}^{\rho_{i,i}-1}$. Then by Lemma 7.15, we have either $t' = r_{\eta_i+2j'}$ with $0 \leq j' \leq \lambda_{i,i} - 2$, or $t' \notin M_i$. Let h be the largest index $0 \leq h < i$ such that $\lambda_{i,h} = \lambda_{i,i} - 1$. Then the maximality of h and the condition 3 for M_i imply that $\lambda_{i,h+1} = \lambda_{i,i}$, ω_h is a wide transformation with support $K_h = \{r_{\eta_i+2\lambda_{i,i}-2+j'}\}_{j'=0}^{N+1}$ and $\lambda_{i,h'} \geq \lambda_{i,i}$ for every $h+1 < h' < i$.

Now we show that $t' \in [y^{(h')}]$ for every $h \leq h' \leq i$. This follows from the choice of h in the case that $t' = r_{\eta_i+2j'}$ with $0 \leq j' \leq \lambda_{i,i} - 2$. Thus we may assume that $t' \notin M_i$. Now we apply the current inductive argument for construction of the sets $M_0, M_1, \dots, M_i$ to $\omega_{i+1}^{-1} \cdots \omega_n^{-1} \omega_0^{-1} \omega_1^{-1} \cdots \omega_i^{-1} \in Y_{y^{(i+1)}, y^{(i+1)}}$ instead of $w = \omega_n \cdots \omega_1 \omega_0 \in Y_{x,x}$. Namely, the sets $J^{(i+1)} = \{r_{\eta_i+2\lambda_{i,i}+j'}\}_{j'=0}^{N-3} \cup \{t, t'\}$ and $J^{(i+1)} \cup \{r_{\eta_i+2\lambda_{i,i}+N-2}, r_{\eta_i+2\lambda_{i,i}+N-1}\} = J^{(i)} \cup \{t, t'\}$ play the roles of M_0 and M_1 , respectively, the sets $J^{(i+1)}$ and $J^{(i)}$ play the roles of $J^{(0)}$ and $J^{(1)}$, respectively, and so on. Now each $\omega_{i+1-h'}^{-1}$ with $1 \leq h' \leq i-h$ (at most i members in total) is not a wide transformation with support containing t' . By applying the argument for construction of $M_0, M_1, \dots, M_{i-h}$ to our present situation, this implies that every $y^{(i+1-h')}$ ($1 \leq h' \leq i-h$) (hence $y^{(h+1)}$, by putting $h' = i-h$) contains t' . Moreover, by the above property of ω_h , t' is not contained in K_h and it is also contained in $y^{(h)}$. Thus the claim of this paragraph holds. Let $\lambda \in \Lambda$ be the index such that $t' = (y^{(h)})_\lambda$.

Now by an argument similar to Lemma 7.14, each $\omega_{h'}$ ($h+1 \leq h' < i$) is one of the following three types:

- A wide transformation with support $K_{h'} \subseteq \{r_{\eta_i+j'} \mid j' \geq 2\lambda_{i,i}\}$.
- A narrow transformation with support $K_{h'}$ containing neither an element of $[y^{(h')}] \cap M_i$ nor t' .
- A narrow transformation of not of type 1 in Section 7.5.1 with support $K_{h'}$ containing either an element of $[y^{(h')}] \cap M_i$ or t' .

Then by an argument similar to Lemmas 7.14 and 7.15 (using Lemmas 7.7 and 7.8), there exist a $z \in S^{(\Lambda)}$ and an element u such that $[z] = [y^{(i)}]$, $u \in Y_{z,y^{(h)}}$, $u \cdot \alpha_{\eta_i+2\lambda_{i,i}-2} = \alpha_{\eta_i+2\lambda_{i,i}}$, u is a product of wide or narrow transformations with support not containing $t' = (y^{(h)})_\lambda$ (hence not containing t), and $u \in \widehat{Y}_{z',y'}$, where y' and z' are obtained from $y^{(h)}$ and z by replacing $(y^{(h)})_\lambda = z_\lambda = t'$ with $r_{\eta_i+2\lambda_{i,i}-2} \in J^{(h)}$ and $r_{\eta_i+2\lambda_{i,i}} \in J^{(i)}$, respectively. Now by the property of the wide transformation ω_i mentioned above, the support $K_i = J^{(i)} \cup \{t, t'\}$ of ω_i is of type A_{N+2} and is not adjacent to $[y^{(i)}] \setminus \{t'\} = [z'] \setminus \{r_{\eta_i+2\lambda_{i,i}}\}$. Since $N \geq 2$, this implies that $\beta = \alpha_t + \alpha_{\eta_i+2\lambda_{i,i}} + \alpha_{\eta_i+2\lambda_{i,i}+1} \in \Pi^{[z']}$, therefore $\gamma = u^{-1} \cdot \beta \in \Pi^{[y']}$. Note that $[y'] = [y^{(i+1)}]$ by the choices of y' and $y^{(i)}$, and the property of ω_i . Thus there is a $z'' \in S^{(\Lambda)}$ such that $[z''] = [z']$ and $u \in Y_{z'',y^{(i+1)}}$, therefore $\gamma \in \Pi^{[y^{(i+1)}]}$. Moreover, by the above property of u , the coefficients of α_t and $\alpha_{t'}$ in γ are equal to those of α_t and $\alpha_{t'}$, respectively, in β ; namely 1 and 0. This implies that $\gamma \in \Pi^{[y^{(i+1)}]} \setminus \Phi_{J^{(i+1)}}$ is not orthogonal to $\alpha_{t'} \in \Pi_{J^{(i+1)}}$. In the same way as Lemma 7.14, this leads a contradiction that $(\omega_i \cdots \omega_1 \omega_0)^{-1} \cdot \gamma \in \Pi^{[x]} \setminus \Pi_L$ is not orthogonal to an element of Π_L . Hence the proof of Lemma 7.16 is concluded. $\square$

Moreover, we have the following result:

Lemma 7.17. *In the above situation, the case $t' \in \{r_{\eta_i+2\lambda_{i,i}+N+1+2j}\}_{j=0}^{\rho_{i,i}-1}$ does not occur.*

Proof. Assume contrary that $t' = r_{\eta_i+2\lambda_{i,i}+N+1+2j_0}$ with $0 \leq j_0 \leq \rho_{i,i} - 1$. Let h be the largest index $0 \leq h \leq i$ such that $r_{\eta_i+2\lambda_{i,h}+N+1} = t'$ (i.e., $\lambda_{i,h} = \lambda_{i,i} + j_0$). Then the maximality of h and the condition 3 for M_i imply that $\lambda_{i,h'} \leq \lambda_{i,i} + j_0$ for every $h \leq h' \leq i$. Let $\lambda \in \Lambda$ be the index such that $t' = (y^{(h)})_\lambda$.

Now by an argument similar to Lemma 7.14, each $\omega_{h'}$ ($h \leq h' < i$) is one of the following three types:

- A wide transformation with support $K_{h'} \subseteq \{r_{\eta_i+j'} \mid j' \leq 2\lambda_{i,i} + N - 1 + 2j_0\}$.
- A narrow transformation with support $K_{h'}$ containing no elements of $[y^{(h')}] \cap M_i$.
- A narrow transformation of not of type 1 in Section 7.5.1 with support $K_{h'}$ containing some element of $[y^{(h')}] \cap M_i$.

Then by an argument similar to Lemmas 7.14, 7.15, and 7.16 (using Lemmas 7.7 and 7.8), there exist a $z \in S^{(\Lambda)}$ and an element u such that $[z] = [y^{(i)}]$, $u \in Y_{z,y^{(h)}}$, $u \cdot \alpha_{\eta_i+2\lambda_{i,i}+2j_0} = \alpha_{\eta_i+2\lambda_{i,i}}$, u is a product of wide or narrow transformations with support not containing $t' = (y^{(h)})_\lambda$ (hence not containing t), and $u \in \widehat{Y}_{z',y'}$, where y' and z' are obtained from $y^{(h)}$ and z by replacing $(y^{(h)})_\lambda = z_\lambda = t'$ with $r_{\eta_i+2\lambda_{i,i}+2j_0} \in J^{(h)}$ and $r_{\eta_i+2\lambda_{i,i}} \in J^{(i)}$, respectively. Now by the property of the wide transformation ω_i mentioned above, the support $K_i = J^{(i)} \cup \{t, t'\}$ of ω_i is of type A_{N+2} and is not adjacent to $[y^{(i)}] \setminus \{t'\} = [z'] \setminus \{r_{\eta_i+2\lambda_{i,i}}\}$. Since $N \geq 2$, this implies that $\beta = \alpha_t + \alpha_{\eta_i+2\lambda_{i,i}} + \alpha_{\eta_i+2\lambda_{i,i}+1} \in \Pi^{[z']}$, therefore $\gamma = u^{-1} \cdot \beta \in \Pi^{[y']}$. Let h_0 be an index such that $\lambda_{i,h_0} = \lambda_{i,i} + j_0 + 1$ (i.e., $t' = r_{\eta_i+2\lambda_{i,h_0}+N-1} \in J^{(h_0)}$). Then by the choices of y' and $y^{(i)}$, we have $[y'] = [y^{(h_0)}]$. Thus there is a $z'' \in S^{(\Lambda)}$ such that $[z''] = [z']$ and $u \in Y_{z'',y^{(h_0)}}$, therefore $\gamma \in \Pi^{[y^{(h_0)}]}$. Moreover, by the above property of u , the coefficients of α_t and $\alpha_{t'}$ in γ are equal

to those of α_t and $\alpha_{t'}$, respectively, in β ; namely 1 and 0. This implies that $\gamma \in \Pi^{[y^{(h_0)}]} \setminus \Phi_{J^{(h_0)}}$ is not orthogonal to $\alpha_{t'} \in \Pi_{J^{(h_0)}}$. In the same way as Lemma 7.14, this leads a contradiction that $(\omega_{h_0-1} \cdots \omega_1 \omega_0)^{-1} \cdot \gamma \in \Pi^{[x]} \setminus \Pi_L$ is not orthogonal to an element of Π_L . Hence the proof of Lemma 7.17 is concluded. $\square$

Hence by Lemmas 7.16 and 7.17, we have $\lambda_{i,i} = 0$ in the case that ω_i is a wide transformation. Put $r_{\eta_i-1} = t$ and $r_{\eta_i-2} = t'$. Now by applying the current inductive argument for construction of the sets $M_0, M_1, \dots, M_i$ to $\omega_{i+1}^{-1} \cdots \omega_n^{-1} \omega_0^{-1} \omega_1^{-1} \cdots \omega_i^{-1} \in Y_{y^{(i+1)}, y^{(i+1)}}$ instead of $w = \omega_n \cdots \omega_1 \omega_0 \in Y_{x,x}$, it follows (in a way similar to the second paragraph in the proof of Lemma 7.16) that $t' = r_{\eta_i-2}$ is contained in every $y^{(h)}$, $1 \leq h \leq i$. Moreover, since $t' \notin M_i$ and we assumed that ω_0 is a wide transformation, t' is not contained in the support of ω_0 , therefore $t' \in [y^{(0)}]$. Now the data defined by $M_{i+1} = M_i \cup \{t, t'\}$, $\eta_{i+1} = \eta_i - 2$, $\mu_{i+1} = \mu_i + 2$, $\lambda_{i+1,h} = \lambda_{i,h} + 1$, $\rho_{i+1,h} = \rho_{i,h}$ ($0 \leq h \leq i$), $\lambda_{i+1,i+1} = 0$, and $\rho_{i+1,i+1} = \rho_{i,i}$ satisfy the above four conditions. Thus we have determined the data for the case of index $i+1$.

By the above argument, it follows inductively that the above data exist for the case of index $i = n+1$, where $y^{(n+1)} = x$. Then by comparing the sets $[y^{(0)}] \cap M_{n+1}$ and $[y^{(n+1)}] \cap M_{n+1}$ in the condition 2, it follows from the fact $y^{(0)} = y^{(n+1)} = x$ that $\lambda_{n+1,n+1} = \lambda_{n+1,0}$ and $\rho_{n+1,n+1} = \rho_{n+1,0}$, therefore $J^{(n+1)} = J^{(0)} = L$. Moreover, by the condition 3 and the property of the wide transformations (see Section 7.5.1), we have $w \cdot \alpha_{\eta_{n+1}+2\lambda_{n+1,0}+j} = \alpha_{\eta_{n+1}+2\lambda_{n+1,0}+j}$ for every $0 \leq j \leq N-1$, therefore w fixes $\Pi_{J^{(0)}} = \Pi_L$ pointwise. This contradicts our assumption.

Hence the claim holds in the case that L is of type A_N , $N \geq 2$.

7.5.3 Proof for the case $N = 1$

In this subsection, we suppose that L is of type A_1 and prove that w fixes Π_L . The argument for the case is analogous to that in Section 7.5.2. Namely, let $w = \omega_n \cdots \omega_1 \omega_0$, where $\omega_i = \omega_{y^{(i)}, J^{(i)}}^{t^{(i)}}$ for each i , be a decomposition of $w \in Y_{x,x}$ as in Section 7.3 with respect to $J^{(0)} = L$. Then in the same way as Section 7.5.2, we may assume without loss of generality that the expressions $w_i = \omega_{i-1} \cdots \omega_1 \omega_0 \omega_n \cdots \omega_i$ and $w_i^{-1} = \omega_i^{-1} \cdots \omega_n^{-1} \omega_0^{-1} \omega_1^{-1} \cdots \omega_{i-1}^{-1}$ are also decompositions of w_i and w_i^{-1} , respectively, as in Section 7.3 (i.e., the length of the product is equal to the sum of the lengths of the factors); and that ω_0 is a wide transformation of type 1 in Section 7.5.1 (note that a wide transformation of type 2 fixes the set $\Pi_{L'}$ pointwise). For each h , let t_h denote the unique element of $J^{(h)}$. From now, we determine inductively subsets M_i of S such that $M_{i-1} \subseteq M_i$ and the following conditions are satisfied:

1. Any pair of elements of M_i is not adjacent to each other in Γ .
2. For each $0 \leq h \leq i$, we have $M_i = ([y^{(h)}] \cap M_i) \cup \{t_h\}$, and $[y^{(h)}] \cap M_i$ and $[y^{(h)}] \setminus M_i$ are not adjacent to each other in Γ .
3. For each $1 \leq h \leq i$,
 - if $t_{h-1} \neq t_h$, then ω_{h-1} is a wide transformation of type 1 that exchanges $\alpha_{t_{h-1}}$ and α_{t_h} and fixes $[y^{(h-1)}] \setminus \{t_h\} = [y^{(h)}] \setminus \{t_{h-1}\}$ pointwise;
 - if $t_{h-1} = t_h$, then ω_{h-1} is either a narrow transformation or a wide transformation of type 2.
4. For any $t \in M_i$, there exists some $0 \leq h \leq i$ such that $t_h = t$.

For the case $i = 0$, we define $M_0 = L = \{t_0\}$, which satisfies the above conditions. Now suppose that $i \leq n$ and we have determined such a subset M_i . Then for the possibility of the next transformation ω_i , we have the following result:

Lemma 7.18. *In the setting, the following situation does not occur: ω_i is a narrow transformation of type 1 among the ten types in Section 7.5.1 whose support K_i contains an element of $[y^{(i)}] \cap M_i$.*

Proof. The proof is analogous to, and much simpler than, the one for Lemma 7.14. Assume contrary that ω_i is a narrow transformation of type 1 whose support K_i contains an element s of $[y^{(i)}] \cap M_i$. Now $\{s\}$ is an irreducible component of $y^{(i)}$. By the definition of narrow transformations of type 1, we have $K_i = \{s, s'\}$ for some $s' \in S \setminus ([y^{(i)}] \cup \{t_i\})$ and $m_{s,s'}$ is odd. Now s' is not adjacent to $([y^{(i)}] \cup \{t_i\}) \setminus \{s\}$. By the condition 4 for M_i , there is an index $0 \leq h < i$ such that $s = t_h$, therefore $[y^{(h)}] = ([y^{(i)}] \cup \{t_i\}) \setminus \{s\}$. This implies that $\alpha_{s'} \in \Pi^{[y^{(h)}]}$ and $\alpha_{s'}$ is not orthogonal to $\alpha_s = \alpha_{t_h}$, therefore $\gamma = (\omega_{h-1} \cdots \omega_1 \omega_0)^{-1} \cdot \alpha_{s'} \in \Pi^{[x]}$ is not orthogonal to $\alpha_{t_0} = (\omega_{h-1} \cdots \omega_1 \omega_0)^{-1} \cdot \alpha_{t_h} \in \Pi_L$. This contradicts the definition of L . Hence Lemma 7.18 holds. $\square$

Now by Lemmas 7.18 and 7.11, if ω_i is a narrow transformation, then either the support of ω_i contains no elements of M_i , or ω_i leaves the set $[y^{(i)}]$ invariant. This implies that the set $M_{i+1} = M_i$ satisfies the above conditions in this case.

From now, we consider the remaining case that ω_i is a wide transformation. If it is of type 2 among the two types in Section 7.5.1, then ω_i fixes $\Pi_{J(i)} = \{\alpha_{t_i}\}$ and leaves the set $[y^{(i)}]$ invariant. On the other hand, suppose that ω_i is a wide transformation of type 1. By the definition, there is an element $s \in [y^{(i)}]$ such that $\{s\}$ is an irreducible component of $y^{(i)}$ and ω_i exchanges α_{t_i} and α_s (hence $s = t_{i+1}$). If $s \in M_i$, then the set $M_{i+1} = M_i$ satisfies the above conditions. On the other hand, suppose that $s \notin M_i$. Then by applying the current inductive argument for construction of the sets $M_0, M_1, \dots, M_i$ to $\omega_{i+1}^{-1} \cdots \omega_n^{-1} \omega_0^{-1} \omega_1^{-1} \cdots \omega_i^{-1} \in Y_{y^{(i+1)}, y^{(i+1)}}$ instead of $w = \omega_n \cdots \omega_1 \omega_0 \in Y_{x,x}$, it follows (in a way similar to the second paragraph in the proof of Lemma 7.16) that s is contained in every $y^{(h)}$, $1 \leq h \leq i$. Moreover, since we assumed that ω_0 is a wide transformation, we have either ω_0 is of type 1 which exchanges α_{t_0} and α_{t_1} , or ω_0 is of type 2 which leaves the sets $[y^{(0)}]$ and $\Pi_{J(0)}$ invariant. Since $s \notin M_i$ and $s \in [y^{(1)}]$, it follows in any case that $s \in [y^{(0)}]$. Thus we have $s \in [y^{(h)}]$ for any $0 \leq h \leq i$, therefore the set $M_{i+1} = M_i \cup \{s\}$ satisfies the above conditions. Hence we have determined the set M_{i+1} .

By the above argument, it follows inductively that such a set M_{n+1} exists, where $y^{(n+1)} = x$. Since $y^{(0)} = x = y^{(n+1)}$, the condition 2 for M_{n+1} implies that $t_{n+1} = t_0$, therefore $w \cdot \alpha_{t_0} = \alpha_{t_{n+1}} = \alpha_{t_0}$ and w fixes $\Pi_L = \{\alpha_{t_0}\}$ pointwise.

Hence the claim holds in the case that L is of type A_1 . Summarizing the results in Sections 7.5.2 and 7.5.3, the claim holds regardless of the type of L . Thus the proof of Theorem 7.1 is concluded.

References

- [1] P. Bahls. *The isomorphism problem in Coxeter groups* (Imperial College Press, 2005).
- [2] A. Björner and F. Brenti. *Combinatorics of Coxeter groups* (Springer, 2005).
- [3] R. E. Borcherds. Coxeter groups, Lorentzian lattices, and K3 surfaces. *Int. Math. Res. Not.* (1998), 1011–1031.
- [4] B. Brink. On centralizers of reflections in Coxeter groups. *Bull. London Math. Soc.* **28** (1996), 465–470.
- [5] B. Brink and R. B. Howlett. Normalizers of parabolic subgroups in Coxeter groups. *Invent. Math.* **136** (1999), 323–351.

- [6] R. Brown. *Topology: A geometric account of general topology, homotopy types and the fundamental groupoid* (Ellis Horwood, 1988).
- [7] D. E. Cohen. *Combinatorial group theory: a topological approach* (Cambridge University Press, 1989).
- [8] V. V. Deodhar. On the root system of a Coxeter group. *Comm. Algebra* **10** (1982), 611–630.
- [9] V. V. Deodhar. Some characterizations of Coxeter groups. *Enseign. Math.* (2) **32** (1986), 111–120.
- [10] V. V. Deodhar. A note on subgroups generated by reflections in Coxeter groups. *Arch. Math.* (Basel) **53** (1989), 543–546.
- [11] M. Dyer. Reflection subgroups of Coxeter systems. *J. Algebra* **135** (1990), 57–73.
- [12] E. Godelle. Normalisateurs et centralisateurs des sous-groupes paraboliques dans les groupes d’Artin-Tits. Ph.D. thesis. University of Picardie-Jules Verne (2001).
- [13] E. Godelle. Normalisateur et groupe d’Artin de type sphérique. *J. Algebra* **269** (2003), 263–274.
- [14] R. B. Howlett. Normalizers of parabolic subgroups of reflection groups. *J. London Math. Soc.* (2) **21** (1980), 62–80.
- [15] R. B. Howlett, P. J. Rowley and D. E. Taylor. On outer automorphism groups of Coxeter groups. *Manuscripta Math.* **93** (1997), 499–513.
- [16] J. E. Humphreys. *Reflection groups and Coxeter groups* (Cambridge University Press, 1990).
- [17] D. L. Johnson. *Presentations of groups, second edition* (Cambridge University Press, 1997).
- [18] K. Nuida. Centralizers of reflections and reflection-independence of Coxeter groups. Preprint (2006). <http://arxiv.org/abs/math.GR/0602165> arXiv:math.GR/0602165.
- [19] K. Nuida. On centralizers of parabolic subgroups in Coxeter groups. Master thesis. University of Tokyo (2003).
- [20] K. Nuida. Almost central involutions in split extensions of Coxeter groups by graph automorphisms. *J. Group Theory* **10** (2007), 139–166.
- [21] K. Nuida. On the direct indecomposability of infinite irreducible Coxeter groups and the Isomorphism Problem of Coxeter groups. *Comm. Algebra* **34** (2006), 2559–2595.
- [22] I. J. Ortiz. The lower algebraic K -theory of Γ_3 . *K-Theory* **32** (2004), 331–355.
- [23] I. J. Ortiz. Erratum: The lower algebraic K -theory of Γ_3 . *K-Theory* **38** (2007), 85–86.
- [24] L. Paris. Centralizers of parabolic subgroups of Artin groups of type A_l , B_l and D_l . *J. Algebra* **196** (1997), 400–435.
- [25] L. Paris. Commensurators of parabolic subgroups of Coxeter groups. *Proc. Amer. Math. Soc.* **125** (1997), 731–738.
- [26] L. Paris. Irreducible Coxeter groups. *Internat. J. Algebra Comput.* **17** (2007), 427–447.

Koji Nuida

Present address: Research Center for Information Security, National Institute of Advanced Industrial Science and Technology, Akihabara-Daibiru Room 1003, 1-18-13 Sotokanda, Chiyoda-ku, Tokyo 101-0021, Japan
E-mail: k.nuida@aist.go.jp